arXiv:1206.3856v1 [math.PR] 18 Jun 2012

Noise-stability and central limit theorems for effective resistance of random electric networks

Raphaël Rossignol

March 8, 2013

Abstract

We investigate the (generalized) Walsh decomposition of point-to-point effective resistances on countable random electric networks with i.i.d resistances. We show that it is concentrated on low levels, and thus point-to-point effective resistances are uniformly stable to noise. For graphs that satisfy some homogeneity property, we show in addition that it is concentrated on sets of small diameter. As a consequence, we compute the right order of the variance and prove a central limit theorem for the effective resistance through the discrete torus of side n in $\mathbb{Z}^d$, when n goes to infinity.

Keywords: Effective resistance, conductance, noise sensitivity, stability, Efron-Stein inequality and decomposition, generalized Walsh decomposition, central limit theorem, stochastic homogenization.

1 Introduction

Consider a piece of conductive material whose resistivity possesses some microscopic disorder. One way to account for this disorder is to suppose that the material is an electric network made of tiny random resistances. Once this model is assumed, one typically wants to understand the behaviour of the macroscopic resistivity of the material. To make the picture more accurate, imagine that each edge of the lattice $\mathbb{Z}^d$ is equipped with a resistance $r(e)$ belonging to some interval $[1,\Lambda]$ (we shall not precise any resistance unit). Suppose in addition that all resistances are random, independently and identically distributed. Our macroscopic piece of material is now the box $\mathcal{B}_n=\{0,\dots,n\}^d$, two sides of which we distinguish: $A_n=\{x\in\mathcal{B}_n \text{ s.t. } x_1=0\}$ and $Z_n=\{x\in\mathcal{B}_n \text{ s.t. } x_1=n\}$. The *effective resistance* of the box $\mathcal{B}_n$ is then defined as:

$$\mathcal{R}_n=\inf_\theta \sum_{e\in\mathsf{E}_n} r(e)\theta^2(e)\ ,$$

where the sum is over the set E_n of the edges inside $\mathcal{B}_n$ and the infimum is taken over all unit flows on E_n from A_n to Z_n (all the precise definitions are postponed until section 2). In the litterature, the *effective conductivity* is more often the main character. It is simply the inverse of the effective resistance and can also be defined as:

$$\mathcal{C}_n=\inf_v \sum_{e\in\mathsf{E}_n} c(e)(dv(e))^2\ ,$$

where the infimum is over all functions v on $\mathcal{B}_n$ having value 0 on A_n and 1 on Z_n, $c(e) = 1/r(e)$ is the conductance of edge e, and $dv(e) := v(e_-) - v(e_+)$ is the difference of v along edge e. The unique minimizer in the definition of $\mathcal{C}_n$ is the function having value 0 on A_n and 1 on Z_n being discrete harmonic on $\mathcal{B}_n \setminus \{A_n, Z_n\}$. It is worth mentioning that the setting above is also relevant to describe the pressure field of a fluid through a weakly porous medium when the circulation of the fluid can be modelled with Poiseuille's law. The central problem is now to understand the asymptotic probabilistic behaviour of $\mathcal{R}_n$ (or, equivalently, of $\mathcal{C}_n$) as n goes to infinity.

The first step in this direction was accomplished in the setting of stochastic homogenization theory (cf. [14], chapter 7). It is shown in [16], section 3, that a law of large numbers holds (see also [22], [17] and [4] for related results). Namely, there is some positive constant μ such that:

$$\frac{1}{n^{d-2}}\mathcal{C}_n \xrightarrow[n\to\infty]{a.s} \mu \quad \text{i.e} \quad n^{d-2}\mathcal{R}_n \xrightarrow[n\to\infty]{a.s} \frac{1}{\mu} \ .$$

To understand the scaling, notice that the function $v_{hom} : x \mapsto x_1/n$ gives an upperbound of order n^{d-2} on the value of $\mathcal{C}_n$, and the Nash-Williams inequality, a standard result in the theory of electrical networks cf. [18], shows that this order is the right one.

A second step is to understand the fluctuations of $\mathcal{C}_n$ and $\mathcal{R}_n$. If the optimal function in the definition of $\mathcal{C}_n$ was v_{hom}, then $\mathcal{C}_n$ would merely be a sum of $\Theta(n^d)$ i.i.d random variables, each of variance $\Theta(n^{-4})$. The variance of $\mathcal{C}_n$ would thus be of order $\Theta(n^{d-4})$, and that of $\mathcal{R}_n$ of order $\Theta(n^{4-3d})$. A lower bound of this order was given by Wehr, cf. [23], under some technical assumptions (see also section 3.2 below). More recently, an upper bound of the same order was obtained by [12] for a different, but closely related quantity. We shall present in more details the work of Gloria and Otto at the end of this introduction.

The main purpose of the present paper is to derive the right order of the variances of $\mathcal{C}_n$ and $\mathcal{R}_n$ and in addition to make a step further in the understanding of their fluctuations by deriving gaussian central limit theorems for these quantities. However, for technical reasons we shall only be able to do this in a translation invariant setting, namely for the effective resistance through the torus, cf. Theorem 5.2. This is the main result of the paper. Our approach to obtain this result is however quite general and not restricted to graphs like $\mathbb{Z}^d$. Indeed, we shall study in section 3 *the generalized Walsh decomposition* of point-to-point effective resistance on general infinite networks. This decomposition, sometimes called the Efron-Stein decomposition, is an extension of the Fourier-Walsh decomposition on the discrete cube and is related to a notion of noise sensitivity introduced in [2]. From now on we shall drop the term "generalized" for simplicity. The Walsh decomposition of a square integrable function f of the resistances reads:

$$f = \sum_S f_S \ ,$$

where the sum runs over all finite subsets of the set of edges, f_S is a function of $(r(e))_{e\in S}$ for any S and f_S is orthogonal to $f_{S'}$ as soon as $S \neq S'$. Our first result, Theorem 3.5, states that the Walsh decomposition of the effective resistance is always concentrated on low levels, i.e on sets of bounded cardinality. It implies that effective resistances are always uniformly stable to noise in the sense of [2], cf. Corollary 3.6, and also that the famous Efron-Stein inequality is always sharp for estimating the variance of the effective resistance, cf. Corollary 3.7. Then, we shall improve this result on a class of graphs which possess some homogeneity property. These graphs, that we shall qualify as having *homogeneous currents* contain all quasi-transitive graphs, cf. Corollary 3.10. On those graphs we shall show that the Walsh decomposition is concentrated on sets of small diameter. This is the key to obtain a central limit theorem since the sets of resistances with bounded diameter exhibit only local dependence, cf. section 4.

In section 5, we shall adapt this general approach to the effective resistance through the discrete torus, deriving the optimal variance estimate and the gaussian central limit theorem already mentioned.

We end this introduction by giving more details on the work [12], and comparing our results to theirs. Their work is close to the homogenization theory framework. Consider the discrete elliptic differential operator $d^*(cd(.))$ corresponding to random, translation invariant and ergodic conductances $c = (c(e))_{e\in\mathbb{E}^d}$ on $\mathbb{Z}^d$. Precise definitions of d^* and d are given in section 2.1, but let us just mention that it gives, for a function v on $\mathbb{Z}^d$:

$$d^*(cdv)(x) = \sum_{y\sim x} c(x,y)(v(x) - v(y)) \ .$$

Then, using the words of [12], homogenization theory (namely [17]) shows that there exists a constant matrix A such that the solution operator of $\div(A\nabla(.))$ describes the large scale behaviour of the solutions operator of $d^*(cd(.))$. Furthermore, A can be characterized by the so called corrector: for any ξ in $\mathbb{R}^d$, there exists a unique function ϕ_ξ on $\mathbb{Z}^d$ (which is a function also of the conductances) such that $\nabla\phi_\xi$ is stationnary, $\phi_\xi(0) = 0$, $\mathbb{E}(\nabla\phi_\xi) = 0$ and such that $g_\xi : x \mapsto \xi.x + \phi_\xi(x)$ is discrete harmonic for $d^*(cd(.))$ on $\mathbb{Z}^d$. Then, A is characterized by:

$$\xi.A\xi = \mathbb{E}\left[\nabla g_\xi.A\nabla g_\xi\right] \ .$$

When the conductances are i.i.d, A equals μ times the identity matrix, and the constant μ is the same as in the law of large numbers of $\mathcal{C}_n$ stated above. We shall fix $\xi = (1, 0, \dots, 0)$ in the sequel. When one is interested in computing μ, Gloria and Otto remark that the preceding characterization is useless. Thus, one has to find a way to efficiently *estimate* μ. The quantity $\mathcal{C}_n/n^{d-2}$ is therefore a reasonable estimator for μ, and this is where the knowledge of its variance, and even of a central limit theorem, may be useful. As Gloria and Otto remark for (a quantity very similar to) $\mathcal{C}_n$ is that it lacks stationnarity, which is "a handicap for error analysis". Next they introduce a stationary approximation of the voltage, namely ϕ_T solving:

$$\frac{1}{T}\phi_T + d^*(c(\xi + d\phi_T)) = 0 \text{ in } \mathbb{Z}^d \ .$$

Let η_L be an averaging cutoff function with support in $(0, n)^d$ (and some extra regularity condition). When T is large with respect ro n, they show that the quantity:

$$\mathcal{A}_n := \sum_{e\in\mathbb{E}^d} c(e)(\xi + d\phi_T)(e)^2\eta_n(e) \ ,$$

is a good proxy for $n^{2-d}\mathcal{C}_n$, and furthermore, they show that the variance of $\mathcal{A}_n$ is of order at most n^{-d}, with some extra polylogarithmic factor in T for $d = 2$. This order coincides with the variance order conjectured above for $n^{2-d}\mathcal{C}_n$.

What we shall obtain in Theorem 5.2 is an optimal variance estimate and a central limit theorem for the effective conductance on the discrete torus of length n when n goes to infinity. To compare our results to those obtained by Gloria and Otto, we shall say that the precise quantity that we analyse is practically computable and stationary. Furthermore, in some sense, the discrete tori converge to $\mathbb{Z}^d$ better than the discrete cubes since they avoid boundary effects. Thus, the effective conductance on the torus may be a better estimator of μ, even if I do not prove in the present paper that it converges (once normalized) to μ. This was not the goal of my paper and will be the purpose of a future work. Secondly, our method works the same way whether $d = 2$ or not, and this is an advantage over Gloria and Otto's result, which makes a distinction between the two. Finally, the fact that we obtain a central limit theorem is really a step further compared to [12] which only obtain variance estimates. On the other side, Gloria and Otto obtain other interesting results, that we do not get by our method, notably concerning the integrability of the corrector itself (Proposition 2.1 in [12]).

2 Preliminaries

2.1 Effective resistance and minimal current

An excellent reference for background on electric networks is the book [18], chapters 2 and 9 and we shall try to stick to its notations.

In the sequel, $\mathsf{G} = (\mathsf{V}, \mathsf{E})$ will be a *countable, oriented, symmetric and connected graph*. Symmetric means that E is a symmetric subset of V^2, i.e each edge of G occurs with both orientations in E, and countable means here that both V and E are countable. When $e \in \mathsf{E}$, we let e_- denote the tail of e and e_+ its head, we denote by $-e := (e_+, e_-)$ the edge e with reversed direction and let $\mathsf{E}_{1/2}$ be a subset of E such that for every edge e, exactly one of e and $-e$ belongs to $\mathsf{E}_{1/2}$.

For every collection $r \in (\mathbb{R}_+^*)^{\mathsf{E}_{1/2}}$, one may define the *electric network* (G, r): it must be understood as a resistive network, where each edge e is a resistor with resistance $r(e)$. We shall sometimes use the notation $c(e)$ to denote the *conductance* of edge e, that is $c(e) = 1/r(e)$. We define *the coboundary operator* d from $\mathbb{R}^{\mathsf{V}}$ to $\mathbb{R}^{\mathsf{E}}$ by:

$$dv(e) = v(e_-) - v(e_+) \ ,$$

and *the boundary operator* d^* from $\mathbb{R}^{\mathsf{E}}$ to $\mathbb{R}^{\mathsf{V}}$ by:

$$d^*\theta(x) = \sum_{e_-=x} \theta(e) \ .$$

For a fixed collection r, we define $\ell^2_-(\mathsf{E}, r)$ as the Hilbert space of *antisymmetric functions on the edges that have bounded energy*:

$$\ell^2_-(\mathsf{E}, r) = \left\{ \theta \in \mathbb{R}^{\mathsf{E}} \text{ s.t. } \theta \sum_{e \in \mathsf{E}_{1/2}} r(e)\theta^2(e) < \infty \text{ and } \forall e \in \mathsf{E},\ \theta(e) = -\theta(-e) \right\} \ ,$$

endowed with the scalar product:

$$(\theta, \theta')_r = \sum_{e \in \mathsf{E}_{1/2}} r(e)\theta(e)\theta'(e) \ .$$

The square of the norm in $\ell^2_-(\mathsf{E}, r)$ of some θ will be denote by

$$\mathcal{E}_r(\theta) := \sum_{e \in \mathsf{E}_{1/2}} r(e)\theta^2(e) \ ,$$

it is called the *energy* of θ. In the sequel, we shall be interested in *elliptic networks*, i.e networks (G, r) for which there is a finite constant $\Lambda \geq 1$ such that $r \in [1, \Lambda]^{\mathsf{E}}$. Of course, all the sets $\ell^2_-(\mathsf{E}, r)$ for $r \in [1, \Lambda]^{\mathsf{E}}$ are the same, and we shall define this common set as $\ell^2_-(\mathsf{E})$:

$$\ell^2_-(\mathsf{E}) := \left\{ \theta \in \mathbb{R}^{\mathsf{E}} \text{ s.t. } \sum_{e \in \mathsf{E}_{1/2}} \theta^2(e) < \infty \text{ and } \forall e \in \mathsf{E},\ \theta(e) = -\theta(-e) \right\} \ .$$

Let I be a non-negative real number, and u and v two vertices of G. A member θ of $\ell^2_-(\mathsf{E}, r)$ is called *a flow of intensity I from u to v* if:

$$d^*\theta = I(\mathbb{1}_u - \mathbb{1}_v) \ .$$

This means θ satisfies the node law on the network, except at u where a net flow of value I enters the network, and at v where a net flow of value I leaves the network. When θ is a flow from u to v, we say that θ is a unit flow if its intensity is 1. We may now define the *effective resistance between two points u and v* on the network (G, r) as the minimal power dissipated by a unit flow between u and v:

$$\mathcal{R}^{u,v}(r) := \inf \left\{ \sum_{e \in \mathsf{E}_{1/2}} r(e)\theta^2(e) \text{ s.t. } \theta \in \ell_-(\mathsf{E}, r) \text{ is a unit flow from } u \text{ to } v \right\} \ .$$

Since one minimizes a norm on a closed convex set, the infimum above is attained by a unique flow which has the additional property of being a *current.* It is called *the minimal unit current from u to v*, cf. Proposition 9.2 of [18]. We shall denote it by $i^{u,v}_r$. The name *current'* refers to a flow satisfying Kirchhoff's laws as precised in the following definition.

Definition 2.1 *For any $e \in \mathsf{E}$, let $\chi^e = \mathbb{1}_{\{e\}} - \mathbb{1}_{\{-e\}}$ denote the unit flow along e. A* current *$i \in \ell^2_-(\mathsf{E}, r)$ from u to v is a flow from u to v which satisfies* Kirchhoff's laws*:*

- The node law*: for every vertex x different from u and v,*

$$d^*i(x) = 0 \ .$$

- The cycle law*: if $e_1, \ldots, e_n$ is an oriented cycle in G, then:*

$$\left(\sum_{i=1}^{n} \chi^{e_k}, i \right)_r = 0 \ .$$

Currents from u to v of prescribed intensity may or may not be unique depending on the particular graph, see chapter 9 in [18]. On finite graphs however, it is well known that currents are unique (see for instance chapter 2 in [18]). In fact, the minimal current on an infinite graph may be obtained as the limit of currents on a sequence of finite graphs. Let us be more precise. Let $(\mathsf{G}_n)_{n\geq 0}$ be a sequence of finite subgraphs of G that *exhausts* G, i.e such that $\mathsf{G}_n \subset \mathsf{G}_{n+1}$ and such that $\mathsf{G} = \bigcup \mathsf{G}_n$. Denote by G_n^W the network obtained from G by identifying all vertices outside G_n as a single vertex. Notice that one may identify the edges of G_n and G_n^W as subsets of E. A *star* is a member of $\ell^2_-(\mathsf{E}, r)$ of the form $\sum_{e_-=x} c(e)\chi_e$ for $x \in \mathsf{V}$. Let $\bigstar$ (resp. $\bigstar_n$) denote the closed subspace spanned by the stars in $\ell^2_-(\mathsf{E}, r)$ (resp. by the stars of G_n^W). A *cycle* is a member of $\ell^2_-(\mathsf{E}, r)$ of the form $\sum_{i=1}^n \chi^{e_k}$ with $e_1, \ldots, e_n$ an oriented cycle in G. Let $\diamondsuit$ (resp. $\diamondsuit_n$) denote the closed subspace spanned by the cycles in $\ell^2_-(\mathsf{E}, r)$ (resp. by the cycles of G_n). For a closed subspace V, denote by P_V the projection on V in $\ell^2_-(\mathsf{E}, r)$. Then, for any finite path γ from u to v, $P_{\bigstar_n}\gamma$ is the unique current on G_n^W between u and v and it converges in $\ell^2_-(\mathsf{E}, r)$ to $i_r^{u,v} = P_\bigstar\gamma$, the minimal current on G from u to v. Also, $P_{\diamondsuit_n^\perp}\gamma$ converges to $i_r^{F,u,v} = P_{\diamondsuit^\perp}\gamma$, the *free current* from u to v. The free current $i_r^{F,u,v}$ is another current, that may or may not be equal to $i_r^{u,v}$ (but has energy no smaller than $i_r^{u,v}$ of course). On $\mathbb{Z}^d$, for instance, it is known that those currents coincides, and this will be useful in section 5. See [18], Propositions 9.1 and 9.2 and section 9.3 for the details.

2.2 Partial derivatives of the effective resistance and the minimal current

The functions $r \mapsto i_r^{u,v}$ and $r \mapsto \mathcal{R}^{u,v}(r)$ are smooth functions, as the next lemma shows. In the sequel, $\partial_e f$ denotes the partial derivative with respect to $r(e)$ of a function f on $(\mathbb{R}_+^*)^{\mathsf{E}_{1/2}}$.

Lemma 2.2 *The functions $r \mapsto i_r^{u,v}(e)$, for any edge e, and $r \mapsto \mathcal{R}^{u,v}(r)$ admit partial derivatives of all order. In addition, for any vertices u, v, and edges e, e',*

(i) $$\forall e' \neq e, \quad \partial_{e'} i_r^{u,v}(e) = \frac{i_r^{u,v}(e')}{r(e')} i_r^{e'}(e) = \frac{i_r^{u,v}(e')}{r(e)} i_r^{e}(e') \ .$$

(ii) $$\forall e, \quad \partial_e i_r^{u,v}(e) = \frac{i_r^{u,v}(e)}{r(e)} (i_r^e(e) - 1) \ .$$

(iii) $$\forall e, \quad \partial_e \mathcal{R}^{u,v}(r) = (i_r^{u,v}(e))^2 \ .$$

Proof: Let us first suppose that G is finite. Then, it is well known that $i_r^{u,v}(e)$ and $\mathcal{R}^{u,v}(r)$ are rational functions of r with no positive pole. See for instance [5], Theorem 2 p. 46. The idea goes back to Kirchhoff (see [15] for an english translation of the original paper). The fact that $\partial_e \mathcal{R}^{u,v}(r) = (i_r^{u,v}(e))^2$ is also known, cf. for instance [18], Exercise 2.69. One easy way to see it is as follows. Let r' be a collection of resistance differing from r only on edge e. Then, using the minimality of $i_r^{u,v}$,

$$\mathcal{E}_{r'}(i_{r'}^{u,v}) - \mathcal{E}_r(i_{r'}^{u,v}) \leq (\mathcal{R}^{u,v}(r') - \mathcal{R}^{u,v}(r)) \leq \mathcal{E}_{r'}(i_r^{u,v}) - \mathcal{E}_r(i_r^{u,v}) \ ,$$

and thus:

$$(r'(e) - r(e))(i_{r'}^{u,v}(e))^2 \leq (\mathcal{R}^{u,v}(r') - \mathcal{R}^{u,v}(r)) \leq (r'(e) - r(e))(i_{r'}^{u,v}(e))^2 \ . \quad (1)$$

Letting $r'(e)$ go to $r(e)$ shows that $\partial_e \mathcal{R}^{u,v}(r) = (i_r^{u,v}(e))^2$.

To compute the partial derivatives of $r \mapsto i_r^{u,v}(e)$, let us differentiate Kirchhoff's laws of Definition 2.1 with respect to $r'(e)$. We obtain:

$$\forall x \in \mathsf{V}, \quad d^* \left[\partial_{e'} i_r^{u,v}\right](x) = 0 \ ,$$

and for every cycle γ on G,

$$\sum_{e \in \gamma} \left[r(e) \partial_{e'} i_r^{u,v}(e) + i_r^{u,v}(e') \chi_{e'}(e)\right] = 0 \ .$$

Thus, if one defines:

$$j(e) = \partial_{e'} i_r^{u,v}(e) + \frac{i_r^{u,v}(e')}{r(e')} \chi_{e'}(e) \ ,$$

we get:

$$\forall x \notin \{e'_-, e'_+\}, \quad d^* j(x) = 0 \ ,$$

and, for every cycle γ on G,

$$\sum_{e \in \gamma} r(e) j(e) = 0 \ .$$

Thus, j is a current between e_- and e_+. Furthermore, its intensity can be deduced from:

$$d^* j(e'_-) = d^* \left[\partial_{e'} i_r^{u,v}\right](e_-) + \frac{i_r^{u,v}(e')}{r(e')} = \frac{i_r^{u,v}(e')}{r(e')}; .$$

Thus, from the unicity of currents on finite graphs, one gets:

$$j = \frac{i_r^{u,v}(e')}{r(e')} i_r^{e'} \ .$$

Consequently,

$$\forall e' \neq e, \quad \partial_{e'} i_r^{u,v}(e) = \frac{i_r^{u,v}(e')}{r(e')} i_r^{e'}(e) \ ,$$

and:

$$\forall e', \quad \partial_{e'} i_r^{u,v}(e') = \frac{i_r^{u,v}(e')}{r(e')} j(e') = \frac{i_r^{u,v}(e')}{r(e')} (i_r^{e'}(e') - 1) \ .$$

Finally, for any $e \neq e'$,

$$\partial^2_{e',e} \mathcal{R}^{u,v}(r) = \partial_{e'} (i_r^{u,v}(e))^2 = 2 i_r^{u,v}(e) \frac{i_r^{u,v}(e')}{r(e')} i_r^{e'}(e) \ ,$$

But since $\partial^2_{e',e} \mathcal{R}^{u,v}(r) = \partial^2_{e,e'} \mathcal{R}^{u,v}(r)$, we obtain:

$$i_r^{u,v}(e) \frac{i_r^{u,v}(e')}{r(e')} i_r^{e'}(e) = i_r^{u,v}(e') \frac{i_r^{u,v}(e)}{r(e)} i_r^{e}(e') \ .$$

Now, take $(u,v) = e$ and $(u,v) = e'$ and notice that $i^e_r(e)$ and $i^{e'}_r(e')$ are always different from zero. We obtain:

$$\frac{i^e_r(e')i^{e'}_r(e)}{r(e')} = \frac{(i^e_r(e'))^2}{r(e)}\ ,$$

and

$$\frac{i^e_r(e')i^{e'}_r(e)}{r(e)} = \frac{(i^{e'}_r(e))^2}{r(e')}\ .$$

Thus, one deduces that $i^e_r(e') = 0$ if and only if $i^{e'}_r(e) = 0$ and in any case,

$$\frac{i^e_r(e')}{r(e)} = \frac{i^{e'}_r(e)}{r(e')}\ . \tag{2}$$

This last relation is called the *reciprocity law*. See [18], chapter 2 for another proof. This concludes the proof of the lemma on finite graphs.

Now, let G be infinite, r belong to $(\mathbb{R}^*_+)^{\mathsf{E}}$ and e, e' in E. As explained in section 2.1, for any u and v, $i^{u,v}_r$ is the limit, in $\ell^2(\mathsf{E}, r)$ of a sequence $i_{r,n}$ of currents on finite graphs. Notably, $i^{u,v}_r(e)$ is the pointwise limit of $i^{u,v}_{r,n}(e)$. From the formulas of the derivatives on finite graphs, one sees that $r(e') \mapsto i^{u,v}_{r,n}(e)$ form an equicontinuous family of functions on any compact interval I of $\mathbb{R}^*_+$. It follows then from Ascoli's Theorem that the convergence of $i^{u,v}_{r,n}(e)$ to $i^{u,v}_r(e)$ is uniform when $r(e')$ runs over I. Then, from the formulas of the derivative $\partial_{e'} i^{u,v}_{r,n}(e)$ one sees that the derivative itself converges uniformly when $r(e')$ runs over I. Then, $r(e') \mapsto i^{u,v}_r(e)$ is differentiable on $\mathbb{R}^*_+$ and its derivative is the limit of the derivatives $\partial_{e'} i^{u,v}_{r,n}(e)$. This shows the formulas for $\partial_{e'} i^{u,v}_{r,n}(e)$. The last formula is then a consequence of (1), since the continuity of $r(e) \mapsto i^{u,v}_r(e)$ has been established. $\square$

Remark 1 *A similar formula was used in [21], Proposition 1 and in [12], Lemma 2.4.*

The formula satisfied by the partial derivatives of $r \mapsto i^{u,v}_r$ in Lemma 2.2 allows us to control $i^{u,v}_r$ after a finite number of modifications of the individual resistances. *In the whole article, $C(\Lambda)$ (resp. $C(\Lambda, \mathsf{G})$) will denote a constant, depending only on Λ (resp. on Λ and G), that may vary from time to time.*

Lemma 2.3 *Let $\Lambda \geq 1$ be some constant, let r and r' belong to $[1,\Lambda]^E$ and for any subset $S \subset \mathsf{E}$, define $r^{S\leftarrow r'}$ by:*

$$r^{S\leftarrow r'}(e) = \begin{cases} r'(e) & \textit{if} \quad e \in S \\ r(e) & \textit{else} \end{cases}$$

Then,

(i) For any $e \in \mathsf{E}$, if $r'(e) \leq r(e)$,

$$|i^{u,v}_r(e)| \leq |i^{u,v}_{r^{e\leftarrow r'}}(e)| \leq \frac{r(e)}{r'(e)} |i^{u,v}_r(e)|$$

(ii) *There is a sequence of finite numbers $(C_n)_{n\geq 1}$, depending only on Λ, such that for any n, any finite subset $S \subset \mathsf{E}$ of size n, any r and r' in $[1,\Lambda]^{\mathsf{E}}$, any edge $e \in S$ and any vertices u and v:*

$$\left(i^{u,v}_{r^{S\leftarrow r'}}(e)\right)^2 \leq C_n \sum_{e'\in S} \left(i^{u,v}_r(e')\right)^2 \ .$$

One may take $C_n = C(\Lambda)^n$.

Proof: First, let us prove *(i)*. Let $e \in \mathsf{E}$ and consider $\{r(e'), e' \neq e\}$ fixed in $(\mathbb{R}_+^*)^{\mathsf{E}}$. To simplify notations, for $x > 0$, define:

$$f(x) := i^{u,v}_{r^{e\leftarrow x}}(e)$$

and

$$g(x) := i^{e}_{r^{e\leftarrow x}}(e) \ .$$

Then one gets from Lemma 2.2:

$$f'(x) = \frac{f(x)}{x}(g(x) - 1)$$

This is a homogeneous differential equation of order 1 on $\mathbb{R}_+^*$ which implies that f is of constant sign: either it is zero on $\mathbb{R}_+^*$, or it is positive on $\mathbb{R}_+^*$, or it is negative on $\mathbb{R}_+^*$. Suppose that it is not identically zero and orient e so that f is positive. Notice that $g(x) \in]0,1]$ for every $x > 0$. Then, f' is negative, which shows that for any $x_0 \leq x_1$,

$$f(x_1) \leq f(x_0) \ .$$

But also,

$$(-\ln f)'(x) \leq \frac{1}{x}$$

and thus:

$$f(x_0) \leq \frac{x_1}{x_0} f(x_1) \ ,$$

This shows *(i)*. It shows also that *(ii)* holds for sets S of size 1.

Let us prove *(ii)* by recurrence on the size of the set S. Suppose that the lemma is true for sets of size not larger than $n \in \mathbb{N}^*$, and let $e \in \mathsf{E}$ and S be a finite subset of edges of size $n+1$ containing e. Let e' be an edge of S different from e and write $S = \{e'\} \cup S'$ with S' of size n, containing e and not e'. Consider $\{r(e''), e'' \neq e'\}$ fixed in $(\mathbb{R}_+^*)^{\mathsf{E}}$. One gets from Lemma 2.2:

$$\partial_{e'} i^{u,v}_r(e) = \frac{i^{u,v}_r(e')}{r(e')} i^{e'}_r(e)$$

Notice that $|i^{e'}_r(e)|$ is not larger than one, and that from the result for sets of size 1,

$$\sup_{x\in[1,\Lambda]} |i^{u,v}_{r^{e'\leftarrow x}}(e')| \leq \Lambda |i^{u,v}_r(e')| \ .$$

Thus, if we define

$$h(x) := i^{u,v}_{r^{e'\leftarrow x}}(e) \ ,$$

we get:

$$\sup_{x\in[1,\Lambda]} |h'(x)| \leq \Lambda |i_r^{u,v}(e')|$$

whence, for every r in $[1,\Lambda]^{\mathsf{E}_{1/2}}$ and x in $[1,\Lambda]$:

$$|i^{u,v}_{r^{e'\leftarrow x}}(e)| \leq |i_r^{u,v}(e)| + \Lambda^2 |i_r^{u,v}(e')| \ .$$

Thus:

$$(i^{u,v}_{r^{e'\leftarrow x}}(e))^2 \leq 2(i_r^{u,v}(e))^2 + 2\Lambda^4 (i_r^{u,v}(e'))^2 \ . \tag{3}$$

Replacing r by $r^{S'\leftarrow r'}$ and x by $r'(e')$ gives:

$$(i^{u,v}_{r^{S\leftarrow r'}}(e))^2 \leq 2(i^{u,v}_{r^{S'\leftarrow r'}}(e))^2 + 2\Lambda^4 (i^{u,v}_{r^{S'\leftarrow r'}}(e'))^2 \ . \tag{4}$$

But also, exchanging e and e' in (3):

$$(i^{u,v}_{r^{e\leftarrow x}}(e'))^2 \leq 2(i_r^{u,v}(e'))^2 + 2\Lambda^4 (i_r^{u,v}(e))^2 \ .$$

Replacing r by $r^{S'\setminus\{e\}\leftarrow r'}$ and x by $r'(e)$ in the last inequality gives:

$$(i^{u,v}_{r^{S'\leftarrow r'}}(e'))^2 \leq 2(i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e'))^2 + 2\Lambda^4 (i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e))^2 \tag{5}$$

Plugging (4) into (5), one gets:

$$(i^{u,v}_{r^{S\leftarrow r'}}(e))^2 \leq 2(i^{u,v}_{r^{S'\leftarrow r'}}(e))^2 + 4\Lambda^4 (i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e'))^2 + 4\Lambda^8 (i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e))^2 \ .$$

Then one may use the induction hypothesis on S' and $S'\setminus\{e\}\cup\{e'\}$ to obtain:

$$(i^{u,v}_{r^{S'\leftarrow r'}}(e))^2 \leq C_n \left(\sum_{e''\in S'} (i_r^{u,v}(e''))^2 \right) \ ,$$

$$(i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e))^2 \leq C_n \left(\sum_{e''\in S'} (i_r^{u,v}(e''))^2 \right)$$

and

$$(i^{u,v}_{r^{S'\setminus\{e\}\leftarrow r'}}(e'))^2 \leq C_n \left(\sum_{e''\in S'\setminus\{e\}\cup\{e'\}} (i_r^{u,v}(e''))^2 \right) \ .$$

Thus,

$$(i^{u,v}_{r^{S\leftarrow r'}}(e))^2 \leq (2 + 4\Lambda^8 + 4\Lambda^4) C_n \left(\sum_{e''\in S} (i_r^{u,v}(e''))^2 \right) \ .$$

This ends the induction step and shows that one may take $C_n = (2+4\Lambda^8+4\Lambda^4)^n$.
□

2.3 The random setting

For any $e \in \mathsf{E}_{1/2}$, we let μ_e denote some probability measure on $\mathbb{R}_+^*$. The collection of resistances, r will be supposed to be random with distribution $\mathbb{P} := \bigotimes_{e\in E_{1/2}} \mu_e$.

We shall always suppose that the resistances are square integrable. Most of the time, we shall in addition suppose that the network is *elliptic* in the sense that there is a constant $\Lambda > 1$ such that $r \in [1,\Lambda]^{\mathsf{E}}$. Finally, we will use the notation:

$$\forall e \in \mathsf{E}, \quad m_p(e) = \mathbb{E}[|r(e) - \mathbb{E}(r(e))|^p]^{1/p} \ .$$

3 The Walsh decomposition

3.1 Definition and basic properties

For any $e \in \mathsf{E}_{1/2}$ let Δ_e be the following operator on $L^2(\mathbb{R}^{\mathsf{E}_{1/2}}, \mathbb{P})$:

$$\Delta_e f(r) = f(r) - \int f(r) \, d\mu_e(r(e)) \, .$$

From now on, $S \subset \mathsf{E}_{1/2}$ will always mean that S is a *finite* subset of $\mathsf{E}_{1/2}$. For f in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}}, \mathbb{P})$ and for any $S \subset \mathsf{E}_{1/2}$, define:

$$f_S(r) = \mathbb{E}\left[\prod_{e \in S} \Delta_e f(r) \middle| r_S \right] = \sum_{T \subset S} (-1)^{|S \setminus T|} \mathbb{E}[f(r) | r_T] \, ,$$

where $r_S = (r(e))_{e \in S}$. Then, $(f_S)_{S \subset \mathsf{E}_{1/2}}$ is an orthogonal decomposition of f known as the Efron-Stein or the (generalized) Walsh decomposition (cf. [6], [11], [13] and [20] for instance). The basic properties of this decomposition are gathered in the following proposition.

Proposition 3.1 *For any f and g in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}}, \mathbb{P})$, if $S \neq S'$, $\mathbb{E}(f_S g_{S'}) = 0$. Furthermore,*

$$f_\emptyset = \mathbb{E}(f) \, ,$$

$$f = \sum_S f_S \, ,$$

for any $e \in \mathsf{E}_{1/2}$,

$$\Delta_e f = \sum_{S \ni e} f_S \, ,$$

and

$$\sum_{e \in \mathsf{E}_{1/2}} \|\Delta_e f\|^2 = \sum_S |S| . \|f_S\|_2^2 \, .$$

The proofs of the results of this section are minor adaptations of well-known facts, so we gathered them in Appendix A. A trivial consequence of Proposition 3.1 is the famous Efron-Stein's inequality (cf. [11]).

Corollary 3.2 (Efron-Stein's inequality) *For any f in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}}, \mathbb{P})$*

$$\mathsf{Var}(f) \leq \sum_{e \in \mathsf{E}_{1/2}} \|\Delta_e f\|^2 \, .$$

It is also clear that the Efron-Stein inequality is an equality if and only if $f = \sum_{|S| \leq 1} f_S$. This means that f is a constant plus a sum of independent random variables. One sees also that if the variance of f is concentrated on functions f_S such that S is small, then the Efron-Stein inequality is essentially sharp.

There is a number of models of statistical physics flavour where the Efron-Stein inequality is not sharp. Chatterjee (cf. [7]) calls this phenomenon “superconcentration”. This holds for instance for the first passage percolation time between two distant points on $\mathbb{Z}^d$ when $d \geq 2$ (cf. [2, 1]), which may be defined in our setting as:

$$T_r(u,v) := \inf_{\gamma : u \to v} \sum_{e \in \gamma} r(e) \ ,$$

where the infimum is over all paths from u to v. Since superconcentration implies that some part of the variance of f is concentrated on large sets, one sees that, informally, it is related to high complexity (or high non-linearity) of the function f. It is also related to some noise-sensitivity of the function. Indeed, there is a close link between the Walsh decomposition and a notion of noise introduced by [2]. Suppose that r and r' are two independent random variables with the same distribution $\mathbb{P} := \bigotimes_{e \in E_{1/2}} \mu_e$. Let $\varepsilon \in]0,1[$. One constructs a noisy version r^ε of r by replacing with probability ε, at random and independently for any edge e, the variable $r(e)$ by $r'(e)$.

Proposition 3.3 *For any $f \in L^2(\mathbb{R}^{\mathsf{E}_{1/2}}, \mathbb{P})$,*

$$\mathbb{E}[f(r^\varepsilon)|r] = \sum_S (1-\varepsilon)^{|S|} f_S(r) \ ,$$

and thus,

$$\mathsf{Cov}(f(r^\varepsilon), f(r)) = \sum_{S \neq \emptyset} (1-\varepsilon)^{|S|} \|f_S\|_2^2 \ ,$$

To see another interpretation of sensitivity to noise, called chaos, see [7]. Contrarily to what happens to the first passage percolation times, we are dealing with a random variable, the effective resistance, which should be extremely immune to noise. Indeed, one is able to adapt very smoothly the optimal flow when resistances are noised. Thus, the Efron-Stein inequality should always be sharp in this context, and this is what we shall prove in the following sections. We shall in fact prove much more in the context of *current-homogeneous graphs*, namely that the Walsh decomposition is concentrated not only on sets of small size, but already on sets of small diameter.

Finally, to end the parallel between first passage percolation and effective resistance, note that there is a mean to interpolate between those two quantities. Indeed, let us define, for $p \in [1,2]$:

$$\mathcal{R}_r^{u,v}(p) = \inf_{\theta : u \to v} \sum_{e \in \gamma} r(e) |\theta(e)|^p \ .$$

Then, $\mathcal{R}_r^{u,v}(1) = T_r(u,v)$ and $\mathcal{R}_r^{u,v}(2) = \mathcal{R}_r^{u,v}$. The quantity $\mathcal{R}_r^{u,v}(p)$ is called the p-resistance between u and v. Since the distribution of the Walsh decomposition is dramatically different when p equals 1 or 2, it would be interesting to investigate the evolution of the Walsh decomposition of $\mathcal{R}_r^{u,v}(p)$ when p varies continuously from 2 to 1.

3.2 Concentration of the Walsh decomposition on low levels

First, let us study the bound given by Efron-Stein inequality.

Lemma 3.4 *For any* $e \in \mathsf{E}_{1/2}$,

$$\frac{m_2(e)^2}{\Lambda^4}\mathbb{E}[(i_r^{u,v}(e))^4] \leq \|\Delta_e \mathcal{R}^{u,v}\|_2^2 \leq \Lambda^4 m_2(e)^2 \mathbb{E}[(i_r^{u,v}(e))^4] \ .$$

Proof: Using Lemma 2.3 and inequality (1),

$$\begin{aligned}
\|\Delta_e \mathcal{R}^{u,v}\|_2^2 &= \mathbb{E}[(\mathcal{R}^{u,v}(r) - \mathcal{R}^{u,v}(r^{e\leftarrow r'}))_+^2] \ , \\
&\geq \mathbb{E}[(r(e) - r'(e))_+^2 (i_r^{u,v}(e))^4] \ , \\
&\geq \frac{1}{\Lambda^2}\mathbb{E}[(r(e) - r'(e))_+^2 (i_{r^{e\leftarrow 1}}^{u,v}(e))^4] \ , \\
&= \frac{m_2(e)^2}{\Lambda^2}\mathbb{E}[(i_{r^{e\leftarrow 1}}^{u,v}(e))^4] \ , \\
&\geq \frac{m_2(e)^2}{\Lambda^4}\mathbb{E}[(i_r^{u,v}(e))^4] \ .
\end{aligned}$$

On the other side,

$$\begin{aligned}
\|\Delta_e \mathcal{R}^{u,v}\|_2^2 &= \mathbb{E}[(\mathcal{R}^{u,v}(r^{e\leftarrow r'}) - \mathcal{R}^{u,v}(r))_+^2] \ , \\
&\leq \mathbb{E}[(r'(e) - r(e))_+^2 (i_r^{u,v}(e))^4] \ , \\
&\leq \Lambda^2 \mathbb{E}[(r'(e) - r(e))_+^2 (i_{r^{e\leftarrow 1}}^{u,v}(e))^4] \ , \\
&= \Lambda^2 m_2(e)^2 \mathbb{E}[(i_{r^{e\leftarrow 1}}^{u,v}(e))^4] \ , \\
&\leq \Lambda^4 m_2(e)^2 \mathbb{E}[(i_r^{u,v}(e))^4] \ .
\end{aligned}$$

□

The following theorem shows that the Walsh decompositions of point-to-point effective resistances are uniformly concentrated (in terms of the L^2-norm) on sets of small size.

Theorem 3.5 *There is a constant* $C(\Lambda)$ *such that for any graph* G *and any pair of vertices* (u, v),

$$\sum_S |S|^2 \|\mathcal{R}_S^{u,v}\|_2^2 \leq C(\Lambda) \sum_{S \neq \emptyset} \|\mathcal{R}_S^{u,v}\|_2^2 \ .$$

Consequently, for any $k \geq 1$, *any graph* G *and any pair of vertices* (u, v)

$$\sum_{|S| \geq k} \|\mathcal{R}_S^{u,v}\|_2^2 \leq \frac{C(\Lambda)}{k^2} \sum_{S \neq \emptyset} \|\mathcal{R}_S^{u,v}\|_2^2 \ .$$

Proof: Proposition 3.1 implies:

$$\sum_S |S|^2 \|\mathcal{R}_S^{u,v}\|_2^2 = \sum_{e,e'} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \ .$$

Now, remark that for $e \neq e'$:

$$\begin{aligned}
\Delta_e \Delta_{e'} \mathcal{R}^{u,v}(r) &= \mathbb{E}\left[\mathcal{R}^{u,v}(r) - \mathcal{R}^{u,v}(r^{e'\leftarrow r'}) - \mathcal{R}^{u,v}(r^{e\leftarrow r'}) + \mathcal{R}^{u,v}(r^{\{e,e'\}\leftarrow r'}) \middle| r\right] \ , \\
&= \mathbb{E}\left[\int_{r'(e)}^{r(e)} \int_{r'(e)}^{r(e)} \partial_{e,e'}^2 \mathcal{R}^{u,v}(r^{(e,e')\leftarrow(x,y)}) \, dxdy \middle| r\right] \ .
\end{aligned}$$

Thus,

$$\|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \leq 4 m_2(e)^2 m_2(e')^2 \mathbb{E}\left[\sup_{(x,y)\in[1,\Lambda]^2} \left(\partial^2_{e,e'} \mathcal{R}^{u,v}(r^{(e,e')\leftarrow(x,y)})\right)^2\right] .$$

We deduce from Lemma 2.2 that:

$$\partial^2_{e,e'} \mathcal{R}^{u,v}(r) = \partial_e [i_r^{u,v}(e')]^2 = 2 i_r^{u,v}(e') i_r^{u,v}(e) \frac{i_r^e(e')}{r(e)} \tag{6}$$

Thus, Lemma 2.3 shows:

$$\|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \leq C(\Lambda) m_2(e)^2 m_2(e')^2 \mathbb{E}\left[(i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2}\right] . \tag{7}$$

Thus,

$$\sum_{e\neq e'} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \leq C(\Lambda) \mathbb{E}\left[\sum_{e\neq e'} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2}\right] .$$

Recall from (2) that

$$\frac{i_r^e(e')}{r(e)} = \frac{i_r^{e'}(e)}{r(e')} .$$

Thus, for any r,

$$(e,e') \mapsto m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2}$$

is a non-negative symmetric function on $\mathsf{E}_{1/2}^2$. This implies that for any $r \in [1,\Lambda]^{\mathsf{E}}$:

$$\begin{aligned}
& \sum_{e\neq e'} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2} \\
\leq\ & 2 \sum_{\substack{(e,e') \text{ s.t.} \\ |i_r^{u,v}(e')| \leq |i_r^{u,v}(e)|}} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2} \\
\leq\ & 2 \sum_{\substack{(e,e') \text{ s.t.} \\ |i_r^{u,v}(e')| \leq |i_r^{u,v}(e)|}} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e))^4 \frac{(i_r^e(e'))^2}{(r(e))^2} \\
\leq\ & 2(\Lambda-1)^2 \sum_e m_2(e)^2 (i_r^{u,v}(e))^4 \sum_{e'} \frac{(i_r^e(e'))^2}{(r(e))^2} \\
\leq\ & 2(\Lambda-1)^2 \sum_e m_2(e)^2 (i_r^{u,v}(e))^4 ,
\end{aligned}$$

where we have used the fact that:

$$\sum_{e'} \frac{(i_r^e(e'))^2}{(r(e))^2} \leq \frac{\mathcal{R}^e(r)}{r(e)^2} \leq \frac{1}{r(e)} \leq 1 .$$

Thus,

$$\sum_{e\neq e'} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \leq C(\Lambda) \sum_e m_2(e)^2 \mathbb{E}[(i_r^{u,v}(e))^4] \; .$$

Thus, using Lemma 3.4,

$$\begin{aligned}
\sum_S |S|^2 . \|\mathcal{R}_S^{u,v}\|_2^2 &= \sum_{e,e'} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 \; , \\
&= \sum_{e\neq e'} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 + \sum_e \|\Delta_e \mathcal{R}^{u,v}\|_2^2 \; , \\
&\leq C(\Lambda) \sum_e \|\Delta_e \mathcal{R}^{u,v}\|_2^2 \; , \\
&= C(\Lambda) \sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \; . \qquad (8)
\end{aligned}$$

Then, for any $k \in \mathbb{N}^*$,

$$\begin{aligned}
\sum_{|S|\geq k} |S| . \|\mathcal{R}_S^{u,v}\|_2^2 &\leq \frac{1}{k} \sum_S |S|^2 . \|\mathcal{R}_S^{u,v}\|_2^2 \; , \\
&\leq \frac{C(\Lambda)}{k} \sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \; ,
\end{aligned}$$

Thus, for every $k \geq C(\Lambda)$,

$$\sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \leq k \sum_{0<|S|<k} \|\mathcal{R}_S^{u,v}\|_2^2 + \frac{1}{2} \sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \; ,$$

whence, for every $k \geq 2C(\Lambda)$,

$$\sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \leq 2k \sum_{0<|S|<k} \|\mathcal{R}_S^{u,v}\|_2^2 \; .$$

Notably,

$$\sum_S |S| . \|\mathcal{R}_S^{u,v}\|_2^2 \leq 2\lceil 2C(\Lambda) \rceil \sum_{S\neq\emptyset} \|\mathcal{R}_S^{u,v}\|_2^2 \; .$$

Plugging this into inequality (8) ends the proof. □

An easy consequence is that point-to-point effective resistances are uniformly stable to noise.

Corollary 3.6 *On any random network with i.i.d resistances, point-to-point effective resistances are uniformly stable to noise:*

$$\inf_{u,v\in \mathsf{V}} \mathsf{Corr}(\mathcal{R}^{u,v}(r), \mathcal{R}^{u,v}(r^\varepsilon)) \xrightarrow[\varepsilon\to 0]{} 1 \; .$$

Proof: Let us fix u and v, and let f denote the function $r \mapsto \mathcal{R}^{u,v}(r)$

$$\begin{aligned}
\mathsf{Cov}(f(r), f(r^\varepsilon)) &= \sum_{S\neq\emptyset} (1-\varepsilon)^{|S|} \|f_S\|^2 \ , \\
&= \mathsf{Var}(f) - \sum_{S\neq\emptyset} (1-(1-\varepsilon)^{|S|}) \|f_S\|^2 \ , \\
&\geq \mathsf{Var}(f) - \log \frac{1}{1-\varepsilon} \sum_{S\neq\emptyset} |S| \|f_S\|^2 \ , \\
&\geq \mathsf{Var}(f) \left(1 - C(\Lambda) \log \frac{1}{1-\varepsilon} \right) \ .
\end{aligned}$$

This shows that $\mathsf{Corr}(\mathcal{R}^{u,v}(r), \mathcal{R}^{u,v}(r^\varepsilon))$ tends uniformly (in u and v) to 1 when ε tends to zero. □

A trivial consequence of Theorem 3.5 is that the Efron-Stein inequality, Corollary 3.2, is always tight for point-to-point effective resistances.

Corollary 3.7 *There is a constant $C(\Lambda)$ such that for any graph G and any pair of vertices (u, v),*

$$\mathsf{Var}(\mathcal{R}^{u,v}) \leq \sum_{e \in \mathsf{E}_{1/2}} \|\Delta_e \mathcal{R}^{u,v}\|_2^2 \leq C(\Lambda) \mathsf{Var}(\mathcal{R}^{u,v}) \ .$$

To finish this section, we shall recall the setting of the effective resistance through a box in $\mathbb{Z}^d$, which was described in the introduction. Let $\mathcal{B}_n = \{0, \dots, n\}^d$ be equipped with random resistances r on its set of edges E_n, and let us define $A_n = \{x \in \mathcal{B}_n \text{ s.t. } x_1 = 0\}$ and $Z_n = \{x \in \mathcal{B}_n \text{ s.t. } x_1 = n\}$. Then, consider the graph with vertex set B_n, where all the vertices of A_n are identified to a single vertex on one side, and all the vertices of Z_n are identified on the other side. The *effective resistance* through the box $\mathcal{B}_n$ is defined as:

$$\mathcal{R}_n = \mathcal{R}_r^{A_n, Z_n} \ .$$

In [23], it is shown that under some hypotheses on the distribution of the resistances,

$$\mathsf{Var}(\mathcal{C}_n) \geq C n^{d-4} \ ,$$

where $\mathcal{C}_n = 1/\mathcal{R}_n$. This bound translates into:

$$\mathsf{Var}(\mathcal{R}_n) \geq C n^{4-3d} \ .$$

Using Corollary 3.7, it is easy to derive a similar bound in our setting. Let us call $i_{r,n}$ the unit current in the definition of $\mathcal{R}_r^{A_n, Z_n}$. Using Lemma 3.4 and

Jensen's inequality twice:

$$\begin{aligned}
\mathsf{Var}(\mathcal{R}_n) &\geq C(\Lambda) \sum_{e \in \mathsf{E}_n} \|\Delta_e \mathcal{R}_n\|_2^2 \ , \\
&\geq C(\Lambda) \sum_{e \in \mathsf{E}_n} m_2(e)^2 \mathbb{E}[i_{r,n}^4(e)] \ , \\
&\geq C(\Lambda) \sum_{e \in \mathsf{E}_n} m_2(e)^2 \mathbb{E}[i_{r,n}^2(e)]^2 \ , \\
&= C(\Lambda) \#(\mathsf{E}_n) \frac{1}{\#(\mathsf{E}_n)} \sum_{e \in \mathsf{E}_n} m_2(e)^2 \mathbb{E}[i_{r,n}^2(e)]^2 \ , \\
&\geq C(\Lambda) \#(\mathsf{E}_n) \left(\frac{1}{\#(\mathsf{E}_n)} \sum_{e \in \mathsf{E}_n} m_2(e) \mathbb{E}[i_{r,n}^2(e)] \right)^2 \ , \\
&\geq C'(\Lambda) \#(\mathsf{E}_n) \left(\frac{1}{\#(\mathsf{E}_n)} \mathbb{E}[\sum_{e \in \mathsf{E}_n} r(e) i_{r,n}^2(e)] \right)^2 \inf_e m_2(e)^2 \ , \\
&= C'(\Lambda) \frac{1}{\#(\mathsf{E}_n)} \mathbb{E}[\mathcal{R}_n]^2 \inf_e m_2(e)^2 \ , \\
&\geq C''(\Lambda) n^{4-3d} \inf_e m_2(e)^2 \ ,
\end{aligned}$$

which is essentially Wehr's lower bound. Notice that our assumption of ellipticity on the resistances is quite strong. Wehr's assumptions are different, but it allows resistances which are not bounded away from zero and infinity. However, it is easy to derive the result above with an assumption weaker than ellipticity, namely some properly chosen moment condition on $r(e)$ and $1/r(e)$.

Finally, let us emphasize the fact that we are unfortunately unable to show that:

$$\mathbb{E}\left[\sum_{e \in \mathsf{E}_n} i_{r,n}^4(e) \right] \leq C.n^{4-3d} \ . \tag{9}$$

Otherwise, we would obtain from Corollary 3.7 the correct order for the variance of $\mathcal{R}_n$. Notice that (9) does not necessarily hold when the resistances are not supposed to have identical distribution. To understand why, let v_r be the discrete harmonic function on $\mathcal{B}_n \setminus (A_n \cup Z_n)$ with value 1 on A_n and 0 on Z_n. Discrete harmonic at x means that:

$$d^*(c.dv_r)(x) = 0 \ .$$

Then, (9) is equivalent to:

$$\mathbb{E}\left[\frac{1}{\# E_n} \sum_{e \in \mathsf{E}_n} (n dv_r(e))^4 \right] \leq C' \ . \tag{10}$$

When n goes to infinity, one may compare v_r to a continuous analog. Let $(c_x)_{x \in \mathbb{R}^d}$ be a deterministic elliptic collection of conductance matrices. Let $\tilde{v}$ be the function on $[0,1]^d$ with value 1 on $\{x \text{ s.t. } x_1 = 0\}$, 0 on $\{x \text{ s.t. } x_1 = 1\}$ and satisfying on $(0,1)^d$:

$$div(c.\nabla \tilde{v}) = 0 \ .$$

The continuous analog of (10) is the fact that $\nabla\tilde{v}$ belongs to $L^4([0,1]^d)$. However, it is well known that this may be false if the ellipticity constant Λ is not close enough to 1. On $\mathbb{R}^2$, a counterexample is given in [10], see the discussion after Proposition 1.1 therein.

3.3 Further results for graphs with homogeneous currents

We shall consider the *graph distance*, denoted by d, on V. Then, if e and e' are two edges in E, let $d(e,e')$ be the maximal distance between two endpoints, one of which is in e and the other in e'. For any edge e and any collection of resistances r in $[1,\Lambda]^{\mathsf{E}_{1/2}}$, the flow i_r^e belongs to $\ell^2_-(\mathsf{E},r)$. Thus,

$$\sum_{e':\ d(e',e)\geq L} r(e')(i_r^e(e'))^2 \xrightarrow[L\to\infty]{} 0\ . \tag{11}$$

Below, we shall be interested in graphs where the above convergence holds *uniformly in e and r*. We shall say that such a graph has *homogeneous currents.*

Definition 3.8 *Let $\mathsf{G}=(\mathsf{V},\mathsf{E})$ be a countable, oriented, symmetric and connected graph. Let $\Lambda\geq 1$ be a real number, and define:*

$$\alpha(\mathsf{G},L,\Lambda)=\sup\left\{\sum_{e':\ d(e',e)\geq L} r(e')(i_r^e(e'))^2\ \text{s.t. } e\in E,\ r\in[1,\Lambda]^E\right\}\ . \tag{12}$$

The graph G is said to have Λ-homogeneous currents *if:*

$$\alpha(\mathsf{G},L,\Lambda)\xrightarrow[L\to+\infty]{} 0\ .$$

It is natural to expect that for every Λ and Λ' strictly larger than 1, G has Λ-homogeneous currents if and only if it has Λ'-homogeneous currents. However, we could not prove this.

The first fundamental observation, due to Mickaël de la Salle, is that for a fixed edge e, the convergence in (11) always hold uniformly in r, thanks to a compacity argument. The reasoning allows also to obtain some regularity of the functions $r\mapsto i_r^{u,v}$ and $r\mapsto\mathcal{R}^{u,v}(r)$ when $[1,\Lambda]^E$ is equipped with the product topology (it will not be used in the sequel).

Proposition 3.9 *Let u and v be two vertices of G, and supose that $(\mathsf{G}_L)_{L\geq 0}$ is a sequence of finite connected graphs that exhausts G and such that G_0 contains u and v. Then,*

(i) $$\sup\left\{\sum_{e'\in\mathsf{G}_L^c} r(e')(i_r^{u,v}(e'))^2\ \text{s.t. } r\in[1,\Lambda]^E\right\}\xrightarrow[L\to+\infty]{} 0\ .$$

(ii) If $[1,\Lambda]^E$ is equipped with the product topology and $\ell^2_-(\mathsf{E})$ with the strong topology, then the following maps are continuous:

$$\left\{\begin{array}{lll}[1,\Lambda]^E & \to & \ell^2_-(\mathsf{E})\\ r & \mapsto & i_r^{u,v}\end{array}\right. \quad \text{and} \quad \left\{\begin{array}{lll}[1,\Lambda]^E & \to & \mathbb{R}\\ r & \mapsto & \mathcal{R}^{u,v}(r)\end{array}\right.$$

Proof: **(due to Mickaël de la Salle)**. Let us fix u and v two vertices of G. We equip $[1,\Lambda]^E$ with the product topology. By Tychonov's theorem, $[1,\Lambda]^E$ is then a compact (metrizable) space. The function $r \mapsto \mathcal{R}^{u,v}(r)$ is an infimum of continuous functions on $[1,\Lambda]^E$, and thus, it is upper semi-continuous on $[1,\Lambda]^E$. Now, define:

$$c := \lim_{L\to+\infty} \sup \left\{ \sum_{e' \in \mathsf{G}_L^c} r(e')(i_r^{u,v}(e'))^2 \text{ s.t. } r \in [1,\Lambda]^E \right\} .$$

One may find a sequence $(r_L)_{L\geq 1}$ in $[1,\Lambda]^E$ such that:

$$c = \lim_{L\to+\infty} \sum_{e' \in \mathsf{G}_L^c} r_L(e')(i_{r_L}^{u,v}(e'))^2 .$$

The sequence $(i_{r_L}^{u,v})_{L\geq 1}$ is bounded in $\ell^2_-(\mathsf{E})$ and $[1,\Lambda]^E$ is compact, thus one may extract a sequence $(r_{L_k})_{k\geq 1}$ converging to some value r in $[1,\Lambda]^E$ and such that $(i_{r_{L_k}}^{u,v})_{k\geq 1}$ converges weakly (by Banach-Alaoglu's theorem) to some $\theta \in \ell^2_-(\mathsf{E})$. Notice that the property of being a unit flow from u to v is closed for the weak topology in $\ell^2_-(\mathsf{E})$. Thus, θ is a unit flow from u to v. Then, from the upper-smicontinuity of $r \mapsto \mathcal{R}^{u,v}(r)$,

$$\begin{aligned}
\mathcal{R}^{u,v}(r) &\geq \limsup_{k\to+\infty} \mathcal{R}^{u,v}(r_{L_k}) \\
&= \limsup_{k\to+\infty} \sum_{e'} r_{L_k}(e')(i_{r_{L_k}}^{u,v}(e'))^2 \\
&= \limsup_{k\to+\infty} \left(\sum_{e' \in \mathsf{G}_{L_k}} r_{L_k}(e')(i_{r_{L_k}}^{u,v}(e'))^2 + \sum_{e' \in \mathsf{G}_{L_k}^c} r_{L_k}(e')(i_{r_{L_k}}^{u,v}(e'))^2 \right) \\
&\geq \limsup_{k\to+\infty} \sum_{e' \in \mathsf{G}_{L_k}} r_{L_k}(e')(i_{r_{L_k}}^{u,v}(e'))^2 + c \\
&\geq \sup_{L'} \limsup_{k\to+\infty} \sum_{e' \in \mathsf{G}_{L'}} r_{L_k}(e')(i_{r_{L_k}}^{u,v}(e'))^2 + c \\
&= \sup_{L'} \sum_{e' \in \mathsf{G}_{L'}} r(e')(\theta(e'))^2 + c \\
&= \sum_{e'} r(e')(\theta(e'))^2 + c \\
&\geq \mathcal{R}^{u,v}(r) + c .
\end{aligned}$$

This shows that $c = 0$ and proves (i).

Now, take any r in $[1,\Lambda]^E$ and some sequence (r_L) converging to r. Again, one may extract a sequence $(r_{L_k})_{k\geq 1}$ such that $(i_{r_{L_k}}^{u,v})_{k\geq 1}$ converges weakly to some $\theta \in \ell^2_-(\mathsf{E})$ which is a unit flow from u to v. Then,

$$\begin{aligned}
\mathcal{R}^{u,v}(r) \geq \limsup_{k\to+\infty} \mathcal{R}^{u,v}(r_{L_k}) \quad &\geq \quad \liminf_{k\to+\infty} \mathcal{R}^{u,v}(r_{L_k}) \\
&\geq \quad \liminf_{k\to+\infty} \sum_{e'\in\mathsf{G}_{L_k}} r_{L_k}(e')(i^{u,v}_{r_{L_k}}(e'))^2 \\
&\geq \quad \sup_{L'} \liminf_{k\to+\infty} \sum_{e'\in\mathsf{G}_{L'}} r_{L_k}(e')(i^{u,v}_{r_{L_k}}(e'))^2 \\
&= \quad \sup_{L'} \sum_{e'\in\mathsf{G}_{L'}} r(e')(\theta(e'))^2 \\
&= \quad \sum_{e'} r(e')(\theta(e'))^2
\end{aligned}$$

Thus:

$$\mathcal{R}^{u,v}(r) \geq \limsup_{k\to+\infty} \mathcal{R}^{u,v}(r_{L_k}) \geq \liminf_{k\to+\infty} \mathcal{R}^{u,v}(r_{L_k}) \geq \sum_{e'} r(e')(\theta(e'))^2 \geq \mathcal{R}^{u,v}(r) \ .$$

Since $i^{u,v}_r$ is the unique minimizer of $\theta \mapsto \sum_{e'} r(e')\theta(e')$ over the unit flows from u to v, this shows at once that $\theta = i^{u,v}_r$ and that $\mathcal{R}^{u,v}(r_{L_k})$ converges to $\mathcal{R}^{u,v}(r)$ as k goes to infinity. Since this is true for any subsequence $(r_{L_k})_k$ such that $(i^{u,v}_{r_{L_k}})_k$ converges weakly, we deduce that $i^{u,v}_{r_L}$ converges weakly to $i^{u,v}_r$ and that $\mathcal{R}^{u,v}(r_L)$ converges to $\mathcal{R}^{u,v}(r)$. Thus, $r \mapsto \mathcal{R}^{u,v}(r)$ is continuous and $r \mapsto i^{u,v}_r$ is continuous if one endows $\ell^2_-(\mathsf{E})$ with the weak topology. Furthermore, recall from section 2.1 that $i^{u,v}_{r_L}$ belongs to $\bigstar$, defined in $\ell^2_-(\mathsf{E}, r_L)$. Since $d^*(i^{u,v}_{r_L} - i^{u,v}_r) = 0$, $i^{u,v}_{r_L} - i^{u,v}_r$ is orthogonal in $\ell^2_-(\mathsf{E}, r)$ to all the stars of $\ell^2_-(\mathsf{E}, r)$. Thus,

$$\|i^{u,v}_{r_L} - i^{u,v}_r\|^2_{r_L} = \sum_{e'\in\mathsf{E}_{1/2}} r_L(e')(i^{u,v}_r(e'))^2 - \mathcal{R}^{u,v}(r_L) \ .$$

From the dominated convergence theorem, since r_L converges pointwise to r,

$$\lim_{L\to\infty} \sum_{e'\in\mathsf{E}_{1/2}} r_L(e')(i^{u,v}_r)^2 = \mathcal{R}^{u,v}(r) \ .$$

Thus,

$$\lim_{L\to\infty} \|i^{u,v}_{r_L} - i^{u,v}_r\|^2_{r_L} = 0 \ ,$$

which shows that $r \mapsto i^{u,v}_r$ is continuous if one endows $\ell^2_-(\mathsf{E})$ with the strong topology. □

Remark 2 *In Proposition 3.9, the ellipticity hypothesis is used in a crucial way, whereas so far one could have imposed some properly chosen moment conditions on $r(e)$ and $1/r(e)$ instead.*

Corollary 3.10 *Let $\mathsf{G} = (\mathsf{V}, \mathsf{E})$ be a countable, oriented, symmetric and connected graph with automorphism group $\mathsf{Aut}(G)$. Suppose that G is* quasi-transitive *in the sense that E is composed of a finite number of distinct orbits under the action of $\mathsf{Aut}(G)$. Then, for any $\Lambda \geq 1$, G has* Λ-homogeneous currents.

Proof: Fix some edge e in E and take G_L to be the graph whose edges are all the edges of E at distance at most L from e and whose vertices are the endpoints of those edges. Then, from Proposition 3.9,

$$\sup\left\{\sum_{e':\, d(e',e)\geq L} r(e')(i_r^e(e'))^2 \text{ s.t. } r\in[1,\Lambda]^E\right\}\xrightarrow[L\to+\infty]{} 0\ .$$

The quasi-transitivity hypothesis implies that there are only a finite number of different values for:

$$\lim_{L\to\infty}\sup\left\{\sum_{e':\, d(e',e)\geq L} r(e')(i_r^e(e'))^2 \text{ s.t. } r\in[1,\Lambda]^E\right\}$$

when e runs overs E. This implies that G has Λ-*homogeneous currents.* □

A consequence of Corollary 3.10 is that $\mathbb{Z}^d$ has homogeneous currents. This can also be seen in a more robust way using the powerful tool of *elliptic Harnack inequality.* Indeed, let $B_L(e)$ be the vertices at distance at most L from e. Then, $\sum_{d(e',e)\geq L} r(e')(i_r^e(e'))^2$ is upper-bounded by the oscillation on $B_L(e)^c$ of the votage induced by the flow i_r^e. Since this voltage is a bounded function, harmonic on $\mathbb{Z}^d\setminus e$, one may then show, using the Harnack inequality of [9] as in [19], section 6, that $\alpha(\mathbb{Z}^d,L,\Lambda)$ decays at least as quickly as a negative power of L. This argument can be carried out on any graph satisfying the conditions of [9] plus the additional condition that the annuli between $B_L(e)$ and $B_{4L}(e)$ are connected and may be covered by a bounded number of balls of radius L. For instance, this shows that any graph roughly isometric to $\mathbb{Z}^d$ has homogeneous currents.

Now, we end this section with an (artificial) example of a graph which does not have homogeneous currents. For any $k\in\mathbb{N}^*$ let T_k be two copies of complete binary trees of depth k glued at the leafs. The result has two roots. Now, to construct our graph (see Figure 1), we start from $\mathbb{N}^*$, with the usual notion of graph on it, and for any $k\in\mathbb{N}^*$, we do the following construction. We add T_k by glueing one of its root on the vertex $x_{2k}:=2k$ of $\mathbb{N}$ and call the remaining root x'_{2k}. Then, we add a copie T'_k of T_k by glueing one of its root on the vertex $x_{2k+1}:=2k+1$ of $\mathbb{N}$ and call the remaining root x'_{2k+1}. Then we join x'_{2k} and x'_{2k+1} by an edge. We denote by G the resulting graph, let $e_{2k}:=(x_{2k},x_{2k+1})$, $e'_{2k}:=(x'_{2k},x'_{2k+1})$ and equip this graph with unit resistances. Now, the resistance between the root and the leafs of a complete binary tree of any depth is at most $1/2$. Thus, $\mathcal{R}_{T_k}^{x_{2k},x'_{2k}}$, the resistance between x_{2k} and x'_{2k} in the graph T_k is at most 1. Similarly, $\mathcal{R}_{T'_k}^{x_{2k+1},x'_{2k+1}}$ is at most 1. Since resistances in series add, one sees that:

$$i_r^{e_{2k}}(e'_{2k})=\frac{r(e_{2k})}{\mathcal{R}_{T_k}^{x_{2k},x'_{2k}}+r(e'_{2k})+\mathcal{R}_{T'_k}^{x_{2k+1},x'_{2k+1}}}\geq\frac{1}{3}\ .$$

On the other side, the distance between e_{2k} and e'_{2k} is $2k+1$. Thus, for any $\Lambda\geq 1$, and $L\geq 1$,

$$\alpha(\mathsf{G},L,\Lambda)\geq\sum_{e':\, d(e',e_{2L})\geq L} r(e)(i_r^{e_{2L}}(e'))^2\geq(i_r^{e_{2L}}(e'_{2L}))^2\geq\frac{1}{9}\ .$$

Whence G does not have homogeneous currents.



Figure 1: An example of a graph without homogeneous currents

3.3.1 Concentration on sets of small diameter

A small variation on the proof of Theorem 3.5 allows to obtain the following result, which shows that on graphs with homogeneous currents, the Walsh decomposition is concentrated on sets of small diameter.

Theorem 3.11 *For any graph* G*, any* $\Lambda \geq 1$ *and any* $L \geq 1$*,*

$$\sum_{diam(S)\geq L} \|\mathcal{R}_S^{u,v}\|_2^2 \leq C(\Lambda)\alpha(G,L,\Lambda)\sum_{S\neq\emptyset}\|\mathcal{R}_S^{u,v}\|_2^2 \, .$$

Proof: Let L be a positive integer. From inequality (7), one gets:

$$\sum_{(e,e'):\, d(e,e')\geq L} \|\Delta_e\Delta_{e'}\mathcal{R}^{u,v}\|_2^2 \leq \mathbb{E}\left[\sum_{d(e,e')\geq L} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2}\right] \, .$$

Since "$d(e,e') \geq L$" is a symmetric relation between e and e', one may argue as in the proof of Theorem 3.5 to obtain:

$$\begin{aligned} &\sum_{(e,e'):\, d(e,e')\geq L} m_2(e)^2 m_2(e')^2 (i_r^{u,v}(e'))^2 (i_r^{u,v}(e))^2 \frac{(i_r^e(e'))^2}{(r(e))^2} \\ \leq\ & (\Lambda-1)^2 \sum_e m_2(e)^2 (i_r^{u,v}(e))^4 \sum_{e':\, d(e',e)\geq L} \frac{(i_r^e(e'))^2}{(r(e))^2} \\ \leq\ & (\Lambda-1)^2 \alpha(G,L,\Lambda) \sum_e m_2(e)^2 (i_r^{u,v}(e))^4 \, . \end{aligned}$$

Then, taking expectation, Lemma 3.4 and Theorem 3.5 lead to:

$$\sum_{d(e,e')\geq L} \|\Delta_e\Delta_{e'}\mathcal{R}^{u,v}\|_2^2 \leq (\Lambda-1)^2\alpha(G,L)\sum_{S\neq\emptyset}\|\mathcal{R}_S^{u,v}\|_2^2 \, .$$

Finally, notice that:

$$\begin{aligned}
\sum_{d(e,e')\geq L} \|\Delta_e \Delta_{e'} \mathcal{R}^{u,v}\|_2^2 &= \sum_{d(e,e')\geq L} \sum_{S\supset e,e'} \|\mathcal{R}_S^{u,v}\|_2^2 \ , \\
&= \sum_{S} \sum_{\substack{e,e'\in S \\ d(e,e')\geq L}} \|\mathcal{R}_S^{u,v}\|_2^2 \ , \\
&\geq \sum_{diam(S)\geq L} \|\mathcal{R}_S^{u,v}\|_2^2 \ .
\end{aligned}$$

□

3.3.2 The first level carries a significant weight

A corollary of Theorem 3.11 is that on graphs with homogeneous currents, the first level of the Walsh decomposition carries a significant part of the L^2-norm of the centered point-to-point resistances.

Corollary 3.12 *Suppose that G has Λ-homogeneous currents and degree bounded by δ. Then, there is a constant $C(\Lambda, \mathsf{G})$ such that:*

$$\sum_e \|\mathcal{R}_{\{e\}}^{u,v}\|_2^2 \leq \mathsf{Var}(\mathcal{R}^{u,v}) = \sum_{S\neq\emptyset} \|\mathcal{R}_S^{u,v}\|_2^2 \leq C(\Lambda, \mathsf{G}) \sum_e \|\mathcal{R}_{\{e\}}^{u,v}\|_2^2$$

Proof: Let us fix u and v two vertices of G, and let us drop the superscript u, v to lighten the notations. Since G has Λ-homogeneous currents, one may find L large enough (depending on G and Λ) so that:

$$\sum_{S\neq\emptyset} \|\mathcal{R}_S\|_2^2 \leq 2 \sum_{\substack{diam(S)\leq L \\ S\neq\emptyset}} \|\mathcal{R}_S\|_2^2 \ .$$

For L a positive integer, using Lemma 3.13,

$$\begin{aligned}
\sum_{\substack{diam(S)\leq L \\ S\neq\emptyset}} \|\mathcal{R}_S\|_2^2 &\leq C(\Lambda, L) \sum_{diam(S)\leq L} \sum_{e\in S} \|\mathcal{R}_{\{e\}}\|_2^2 \ , \\
&= C(\Lambda, L) \sum_{e\in \mathsf{E}} \|\mathcal{R}_{\{e\}}\|_2^2 \#\{S \text{ s.t. } e\in S \text{ and } diam(S)\leq L\} \ , \\
&\leq C'(\Lambda, L, \delta) \sum_{e\in \mathsf{E}} \|\mathcal{R}_{\{e\}}\|_2^2 \ .
\end{aligned}$$

□

Lemma 3.13 *For any $p \geq 1$,*

$$C_1(\Lambda, p) m_p(e) \|(i_r^{u,v}(e))^2\|_1 \leq \|\mathcal{R}_{\{e\}}^{u,v}\|_p \leq C_2(\Lambda) m_p(e) \|(i_r^{u,v}(e))^2\|_1 \ ,$$

where:

$$m_p(e) = \left(\mathbb{E}\,|r(e) - \mathbb{E}[r(e)]|^p\right)^{1/p} \ .$$

For any S such that $0 < |S| \le L$,

$$\|\mathcal{R}_S^{u,v}\|_p \le C(\Lambda, L) \sum_{e\in S} \|\mathcal{R}_{\{e\}}^{u,v}\|_p \ .$$

Proof: For any function F in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$ and any $p \ge 1$.

$$\begin{aligned}
\|F_{\{e\}}\|_p^p &= \mathbb{E}\left[|\mathbb{E}(\Delta_e F | r(e))|^p\right] \ , \\
&= \mathbb{E}\left[\left|\mathbb{E}(F(r^{e\leftarrow r'}) - F(r) | r'(e))\right|^p\right] \ , \\
&\le \mathbb{E}\left[\left|\mathbb{E}(F(r^{e\leftarrow r'}) - F(r) | r(e), r'(e))\right|^p\right] \ , \\
&= 2\mathbb{E}\left[\left|\mathbb{E}(F(r^{e\leftarrow r'}) - F(r) | r(e), r'(e))\mathbb{1}_{r'(e)>r(e)}\right|^p\right] \ .
\end{aligned}$$

Now, recall from (1) that:

$$(\mathcal{R}(r^{e\leftarrow r'}) - \mathcal{R}(r))\mathbb{1}_{r'(e)>r(e)} \le (r'(e) - r(e))_+ i_r^2(e) \ .$$

Thus, using Lemma 2.3,

$$\begin{aligned}
\|\mathcal{R}_{\{e\}}\|_p^p &\le 2\mathbb{E}[(i_r^2(e)(r'(e) - r(e))_+)^p] \ , \\
&\le C(\Lambda)^p m_p(e)^p \mathbb{E}[(\mathbb{E}(i_r^2(e)|r(e))^p] \ , \\
&\le C'(\Lambda)^p m_p(e)^p \mathbb{E}(i_r^2(e))^p \ .
\end{aligned}$$

On the other side, for any function F in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$ and any $p \ge 1$

$$\begin{aligned}
\|F_{\{e\}}\|_p^p &= \mathbb{E}\left[\left|\mathbb{E}(F(r^{e\leftarrow r'})|r'(e)) - \mathbb{E}(F(r))\right|^p\right] \ , \\
&\ge \frac{1}{2^p}\mathbb{E}\left[\left|\mathbb{E}(F(r^{e\leftarrow r'})|r'(e)) - \mathbb{E}(F(r)|r(e))\right|^p\right] \ , \\
&= \frac{1}{2^{p-1}}\mathbb{E}\left[\left(\mathbb{E}(F(r^{e\leftarrow r'})|r'(e)) - \mathbb{E}(F(r)|r(e))\right)_+^p\right] \ .
\end{aligned}$$

Now, using (1) and Lemma 2.3,

$$\begin{aligned}
\|\mathcal{R}_{\{e\}}\|_p^p &\ge \frac{1}{2^{p-1}}\mathbb{E}\left[\left(\mathbb{E}((r'(e) - r(e))_+ i_r^2(e)|r(e), r'(e))\right)^p\right] \ , \\
&\ge m_p(e)^p C_1(\Lambda, p)\mathbb{E}\left[\left(\mathbb{E}(i_r^2(e)|r(e))\right)^p\right] \ , \\
&\ge m_p(e)^p C_1'(\Lambda, p)\left(\mathbb{E}(i_r^2(e))\right)^p \ .
\end{aligned}$$

Finally, let S be such that $0 < |S| \le L$ and let $e \in S$. For a subset $I \subset E$, let L_I be the operator on $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$ defined by:

$$L_I f(r) = \int f(r) \prod_{e\in I} d\mu_e(r(e)) \ .$$

Notice first that for any function f in $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$,

$$\begin{aligned}
f_S(r) &= \mathbb{E}\left[\left(\prod_{e'\in S\setminus\{e\}}(1-L_{\{e\}})\right)\Delta_e f(r)\middle| r_S\right] ,\\
&= \mathbb{E}\left[\left(\sum_{I\subset S\setminus\{e\}}(-1)^{|I|}L_I\right)\Delta_e f(r)\middle| r_S\right] ,\\
&= \mathbb{E}\left[\sum_{I\subset S\setminus\{e\}}(-1)^{|I|}L_I(\Delta_e f)(r)\middle| r_S\right] .
\end{aligned}$$

Thus, using Jensen's inequality,

$$\|f_S\|_p^p \leq 2^{p|S\setminus\{e\}|}\mathbb{E}\left[\mathbb{E}\left(|\Delta_e f(r)|\,|\,r_S\right)^p\right] .$$

Using (1) and Lemma 2.3,

$$\begin{aligned}
\mathbb{E}\left[\mathbb{E}\left(|\Delta_e\mathcal{R}(r)|\,|\,r_S\right)^p\right] &\leq 2^p\mathbb{E}\left[\mathbb{E}\left((\mathcal{R}(r^{e\leftarrow r'})-\mathcal{R}(r))_+\middle| r_S\right)^p\right] ,\\
&\leq C(\Lambda)^p m_p(e)^p\mathbb{E}\left[\mathbb{E}\left(i_r^2(e)\,|\,r_S\right)^p\right] ,\\
&\leq C(\Lambda)^{pL}m_p(e)^p\left(\sum_{e'\in S}\mathbb{E}\left[i_r^2(e')\right]\right)^p .
\end{aligned}$$

Since this is true for any $e\in S$, we have:

$$\begin{aligned}
\|\mathcal{R}_S\|_p &\leq C(\Lambda)^L\min_{e\in S}m_p(e)\sum_{e'\in S}\mathbb{E}\left[i_r^2(e')\right] ,\\
&\leq C(\Lambda)^L\sum_{e'\in S}m_p(e')\mathbb{E}\left[i_r^2(e')\right] ,\\
&\leq C'(\Lambda,L)\sum_{e'\in S}\|\mathcal{R}^{u,v}_{\{e'\}}\|_p .
\end{aligned}$$

□

4 Central limit theorem

Even if one takes two vertices u and v far appart, there is not necessarily a gaussian central limit theorem for the effective resistance between them, since the influence of an edge near u, for instance, may well represent a positive fraction of the total variance of the resistance. However, let us define the *influence of an edge* e on the effective resistance between u and v by:

$$I^{u,v}(e) := \|\mathcal{R}^{u,v}_{\{e\}}\|_2^2 .$$

Then, if the maximal influence of an edge is small with respect to the variance and if the graph has homogeneous currents and bounded degree, one may obtain a gaussian approximation. The following theorem shows a result in this direction, whereas another instance of this phenomenon will be described on a sequence of finite graphs, the discrete tori, in section 5.

Theorem 4.1 *Let* G *be a graph with homogeneous currents and bounded degree. For vertices* u *and* v *such that* $\mathsf{Var}(\mathcal{R}^{u,v}) > 0$*, define:*

$$\beta(u,v) = \frac{\sup_{e\in\mathsf{E}} I^{u,v}(e)}{\mathsf{Var}(\mathcal{R}^{u,v})} \ ,$$

and

$$\overline{\mathcal{R}}^{u,v} := [\mathcal{R}^{u,v} - \mathbb{E}(\mathcal{R}^{u,v})]/\sqrt{\mathsf{Var}(\mathcal{R}^{u,v})} \ .$$

Let Φ *be the standard gaussian distribution function and let* $F^{u,v}$ *be the distribution function of* $\overline{\mathcal{R}}^{u,v}$*. There is a function* $f : \mathbb{R}^+ \to [0,1]$*, depending on* G *only, such that:*

$$f(x) \xrightarrow[x\to 0]{} 0 \ ,$$

and:

$$\|F^{u,v} - \Phi\|_\infty \leq f(\beta(u,v)) \ .$$

Proof: Let us fix the vertices u and v. For every integer L, one defines $J = J(L)$ as:

$$J(L) = \{S \subset E \text{ s.t. } diam(S) \leq L \text{ and } S \neq \emptyset\} \ ,$$

Let W_L be the random variable:

$$W_L = \frac{\sum_{S\in J(L)} \mathcal{R}_S^{u,v}}{\sqrt{\sum_{S\in J(L)} \|\mathcal{R}_S^{u,v}\|_2^2}} \ .$$

Since G has homogeneous currents, we know that for L large enough, W_L will be close (in L^2-norm) to $\overline{\mathcal{R}}^{u,v}$. On the other hand, since $\mathcal{R}_S^{u,v}$ depends only on $(r(e))_{e\in S}$, we know that for every fixed L, W_L is a sum of random variables with only local dependence. Thus, one may use the work of [8] to control the distance to normality.

To be more precise, let F_L be the distribution function of W_L. Using the notations of [8], it is easy to see that $N(C_S)$ is included in the subsets of $J(L)$ that lie at distance at most $7L$ from S. Thus, one may use Theorem 2.4 of [8] with $p = 3$ and $\kappa \leq \delta^{8L^2}$ where δ is a bound on the degrees in G. We obtain:

$$\|F_L - \Phi\|_\infty \leq \delta^{8L^2} \frac{\sum_{S\in J(L)} \|\mathcal{R}_S^{u,v}\|_3^3}{\left(\sum_{S\in J(L)} \|\mathcal{R}_S^{u,v}\|_2^2\right)^{3/2}} \ .$$

Using Corollary 3.12 and Lemma 3.13,

$$\begin{aligned}
\|F_L - \Phi\|_\infty &\leq C_1(L,\Lambda,\mathsf{G}) \frac{\sum_{S\in J(L)} \left(\sum_{e\in S} \|\mathcal{R}_{\{e\}}^{u,v}\|_3\right)^3}{\left(\sum_{e\in\mathsf{E}} \|\mathcal{R}_{\{e\}}^{u,v}\|_2^2\right)^{3/2}} \ , \\
&\leq C_1(L,\Lambda,\mathsf{G}) \frac{\sum_{S\in J(L)} |S|^{2/3} \sum_{e\in S} \|\mathcal{R}_{\{e\}}^{u,v}\|_3^3}{\left(\sum_{e\in\mathsf{E}} \|\mathcal{R}_{\{e\}}^{u,v}\|_2^2\right)^{3/2}} \ , \\
&\leq C_1'(L,\Lambda,\mathsf{G}) \frac{\sum_{e\in\mathsf{E}} \|\mathcal{R}_{\{e\}}^{u,v}\|_2^3}{\left(\sum_{e\in\mathsf{E}} \|\mathcal{R}_{\{e\}}^{u,v}\|_2^2\right)^{3/2}} \ ,
\end{aligned}$$

one gets:

$$\|F_L - \Phi\|_\infty \leq C(L)\sqrt{\beta(u,v)} \ ,$$

where $C(L) = C(\Lambda, \mathsf{G}, L)$ is a positive non-decreasing function of L.

On the other hand, Theorem 3.11 ensures that:

$$\|\mathcal{R}^{u,v} - \mathbb{E}(\mathcal{R}^{u,v}) - \sum_{S \in J(L)} \mathcal{R}_S^{u,v}\|^2 \leq \varepsilon(L)\mathsf{Var}(\mathcal{R}^{u,v}) \ ,$$

where $\varepsilon(L) = \varepsilon(\Lambda, \mathsf{G}, L)$ is a positive non-increasing function of L which goes to zero as L goes to infinity. This implies:

$$\|\overline{\mathcal{R}}^{u,v} - W_L\|^2 \leq 4\varepsilon(L) \ .$$

Notice that Φ is 1-Lipschitz (in fact, $1/\sqrt{2\pi}$-Lipschitz), so for any $\eta > 0$ and any $t \in \mathbb{R}$

$$\begin{array}{rcl} F^{u,v}(t) - \Phi(t) & \leq & F^{u,v}(t) - \Phi(t+\eta) + \eta \ , \\ & \leq & F^{u,v}(t) - F_L(t+\eta) + C(L)\sqrt{\beta(u,v)} + \eta \ , \\ & \leq & \mathbb{P}(W_L - \overline{\mathcal{R}}^{u,v} > \eta) + C(L)\sqrt{\beta(u,v)} + \eta \ , \\ & \leq & \dfrac{\|\overline{\mathcal{R}}^{u,v} - W_L\|_2^2}{\eta^2} + C(L)\sqrt{\beta(u,v)} + \eta \ , \\ & \leq & \dfrac{4\varepsilon(L)}{\eta^2} + C(L)\sqrt{\beta(u,v)} + \eta \ . \end{array}$$

Symmetrically, one gets, for any $\eta > 0$ and any $t \in \mathbb{R}$:

$$|F^{u,v}(t) - \Phi(t)| \leq \frac{4\varepsilon(L)}{\eta^2} + C(L)\sqrt{\beta(u,v)} + \eta \ .$$

Optimizing in η gives:

$$\|F^{u,v} - \Phi\|_\infty \leq 3\varepsilon^{1/3}(L) + C(L)\sqrt{\beta(u,v)} \ .$$

It remains to optimize in L. Let $L_0(x) = \sup\{L \in \mathbb{N} \text{ s.t. } 3\varepsilon^{1/3}(L) \geq \sqrt{x}C(L)\}$. Then,

$$\|F^{u,v} - \Phi\|_\infty \leq 6\varepsilon^{1/3}(L_0(\beta(u,v))) \ .$$

Since $L_0(x)$ goes to infinity as x goes to zero, $f(x) := 6\varepsilon^{1/3}(L_0(x))$ answers the Theorem. □

It is in general difficult to apply this result because it is difficult to bound $\beta(u,v)$. However, notice that the influence of an edge is always bounded. Thus, on a bounded graph with homogeneous currents, if the variance of $\mathcal{R}^{u,v}$ goes to infinity as u and v move apart, one gets a central limit theorem. Notice that the last point is equivalent to showing that $\mathbb{E}[\sum_e (i_r^{u,v}(e))^4]$ goes to infinity. It may be shown for instance that this is true on some wedges of $\mathbb{Z}^2$, using the idea of Nash-Williams inequality. For instance, let $h(x) = x^\alpha$ with $\alpha \leq 1/3$, let $\mathsf{V} = \{(x,y) \in \mathbb{Z}^2 \text{ s.t. } |y| \leq h(|x|)$ and let G be the subgraph of $\mathbb{Z}^2$ induced by V. Then, one may derive a central limit theorem for $\mathcal{R}^{0,v}$ on G when the distance $d(0,v)$ goes to infinity. Since this example is not quasi-transitive, one has to use the Harnack inequality to prove that the graph has homogeneous currents.

Notice also that already on $\mathbb{Z}^2$, the variance of the resistance is only of order 1 (cf. [21]), and thus one cannot expect a central limit theorem for point-to-point effective resistance when the resistances are i.i.d (since the influence of the edges near the source and the sink is of order 1). In this respect, the interest of Theorem 4.1 is rather limited. However, one should rather think of it as a first step, with a clean statement, towards central limit theorems for resistances on sequences of finite graphs, as will be made clear in section 5.

Remark 3 (Other divergence form equations) *Let $\mathcal{K}$ be the subset of $\ell^2_-(\mathsf{E})$ consisting of functions of compact support. Define, for any $\theta_0 \in \mathcal{K}$, the following energy:*

$$\mathcal{R}^{\theta_0}(r) := \inf\{ \sum_{e \in \mathsf{E}_{1/2}} r(e)\theta^2(e) \ \text{s.t.}\ \theta \in \ell^2_-(\mathsf{E}) \text{ and } d^*\theta = d^*\theta_0\} \ .$$

When $u, v \in \mathsf{V}$ and θ_0 is a fixed finite path joining u to v, then $\mathcal{R}^{\theta_0}(r) = \mathcal{R}^{u,v}(r)$. It is easy to see that all the results of sections 3 and 4 extend to $(\mathcal{R}^{\theta_0}(r))$, uniformly in $\theta_0 \in \mathcal{K}$.

5 CLT for the effective resistance of the d-dimensional torus

In this section, we investigate when n becomes large the effective resistance of the torus $\mathbb{T}^d_n := (\mathsf{V}^d_n, \mathsf{E}^d_n)$ equipped with non-constant i.i.d resistances on $[1, \Lambda]$. Here, $\mathsf{V}^d_n = (\mathbb{Z}/n\mathbb{Z})^d$ and E^d_n is the traditional set of edges of the torus:

$$\mathsf{E}^d_n = \{(x, y) \in \mathsf{V}^d_n \times \mathsf{V}^d_n \text{ s.t. } \exists i \in \{1, \dots, n\}\ x_i - y_i \in \{-1, 1\} \text{ and } \forall j \neq i,\ x_j = y_j\} \ .$$

One chooses also exactly one edge of each orientation as follows:

$$\mathsf{E}^n_{1/2} = \{(x, y) \in \mathsf{V}^d_n \times \mathsf{V}^d_n \text{ s.t. } \exists i \in \{1, \dots, n\}\ y_i - x_i = 1 \text{ and } \forall j \neq i,\ x_j = y_j\} \ .$$

Since $\mathbb{T}^d_n$ has no natural border, our first objective is to define the effective resistance in a natural and translation invariant way. First, we define a special *cut along direction 1* (see Figure 2):

$$E_0 := \{(x, y) \in \mathsf{E}^n_{1/2} \text{ s.t. } x \sim y,\ x_1 = 0 \text{ and } y_1 = 1\} \ ,$$

and the flows which *cross the torus along direction 1, with intensity 1*:

$$\Theta^n_0 := \{\theta \in \ell^2_-(\mathsf{E}^n_{1/2}) \text{ s.t. } d^*\theta = 0 \text{ and } \sum_{e \in E_0} \theta(e) = 1\} \ .$$

Notice that the elements of Θ^n_0 are sourceless flows and that the definition is translation invariant. Indeed, if we define, for any i in $\{0, \dots, n-1\}$:

$$E_i := \{(x, y) \in \mathsf{V}^d_n \times \mathsf{V}^d_n \text{ s.t. } x \sim y,\ x_1 = i \text{ and } y_1 = i + 1\} \ ,$$

then, for any i and any $\theta \in \ell^2_-(\mathsf{E}^n_{1/2})$, we have:

$$\sum_{e \in E_i} \theta(e) - \sum_{e \in E_{i-1}} \theta(e) = \sum_{x \text{ s.t. } x_1 = i} d^*\theta(x) \ . \qquad (13)$$



Figure 2: The cut E_0 on a 2-dimensional torus.

Thus, for any sourceless flow θ, $\sum_{e \in E_0} \theta(e)$ equals 1 if and only if $\sum_{e \in E_i} \theta(e) = 1$ for some i in $\{0, \dots, n-1\}$. Thus, for any translation τ on the torus,

$$\theta \in \Theta_0^n \Leftrightarrow \theta \circ \tau \in \Theta_0^n \ . \tag{14}$$

Definition 5.1 *One defines* the effective resistance of the torus *as:*

$$\mathcal{R}_n := \inf_{\theta \in \Theta_0^n} \{ \sum_e r(e) \theta^2(e) \} \ . \tag{15}$$

Using the Nash-Williams inequality, it is easy to show that $\mathcal{R}_n = \Theta(1/n^{d-2})$. Using the ideas of the preceding sections, one may show that $\mathcal{R}_n$ satisfies a central limit theorem. This is the main result of the present paper.

Theorem 5.2 *Suppose that* $r(e)$, $e \in \mathsf{E}_{1/2}^n$ *are i.i.d with positive variance and support in* $[1, \Lambda]$ *for some* $\Lambda > 1$. *Then,*

$$\mathsf{Var}(\mathcal{R}_n) = \Theta(n^{4-3d}) \ ,$$

and the resistance satisfies a central limit theorem:

$$\frac{\mathcal{R}_n - \mathbb{E}(\mathcal{R}_n)}{\sqrt{\mathsf{Var}(\mathcal{R}_n)}} \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}(0,1) \ .$$

It is straightforward to show that a similar statement holds for what should be called *the effective conductance of the torus* $\mathbb{T}_n^d$, $\mathcal{C}_n := 1/\mathcal{R}_n$ or *the mean conductivity of the torus* $\mathbb{T}_n^d$: $A_n = n^{2-d} \mathcal{C}_n$. One obtains:

$$\mathsf{Var}(A_n) = \Theta(n^{-d}) \ ,$$

and:

$$\frac{A_n - \mathbb{E}(A_n)}{\sqrt{\mathsf{Var}(A_n)}} \xrightarrow[n \to \infty]{\mathcal{L}} \mathcal{N}(0,1) \ .$$

In [3], the definition of A_n is different, based on the discrete cube and not on the torus, however the behaviour should be the same. [3] obtains only suboptimal bounds for the variance of the mean conductivity. In [12], an optimal bound on the variance is obtained for a related quantity based on the *corrector* of homogenization theory. Again, the behaviour should be the same. In any case, the central limit theorem is new.

Our strategy to show Theorem 5.2 is simply to apply the ideas of Theorem 4.1 to this setting, where the infinite graph is traded against a growing sequence of finite graphs. First, notice that there is a unique minimal flow reaching the infimum in the definition of the resistance.

Lemma 5.3 *The infimum in (15) is attained at a unique flow $i_{r,n}^{per}$. Define the following tangent vector space to Θ_0^n:*

$$\overrightarrow{\Theta} := \{\theta \in \ell^2_-(\mathsf{E}_{1/2}) \text{ s.t. } d^*\theta = 0 \text{ and } \sum_{e \in E_0} \theta(e) = 0\} \ .$$

Then, $i_{r,n}^{per}$ is caracterized (among flows in Θ_0^n) by the following pseudo-Kirchhoff cycle laws:

$$\forall h \in \overrightarrow{\Theta}, \quad (h, i_{r,n}^{per})_r = 0 \ .$$

Proof: Since the function $\theta \mapsto \mathcal{E}_r(\theta)$ is a norm and Θ_0^n is closed and convex, the infimum is unique and attained at some flow $i_{r,n}^{per}$. Then, for any h in $\overrightarrow{\Theta}$ and any $\varepsilon \in \mathbb{R}$, $i_{r,n}^{per} + \varepsilon h$ belongs to Θ_0^n and:

$$\mathcal{E}_r(i_{r,n}^{per} + \varepsilon h) = \mathcal{E}_r(i_{r,n}^{per}) + 2\varepsilon \sum_e r(e)h(e)i_{r,n}^{per}(e) + \varepsilon^2 \mathcal{E}_r(h) \ .$$

Thus, letting ε go to zero, one sees that:

$$\forall h \in \overrightarrow{\Theta}, \quad (h, i_{r,n}^{per})_r = 0 \ .$$

And conversely this condition is enough to caracterize $i_{r,n}^{per}$. It shows also that $i_{r,n}^{per}$ is unique since for any $\theta \in \Theta_0^n$, $\theta - i_{r,n}^{per} \in \overrightarrow{\Theta}$ and:

$$\mathcal{E}_r(\theta) = \mathcal{E}_r(i_{r,n}^{per}) + \mathcal{E}_r(\theta - i_{r,n}) \ .$$

□

Since the minimal flow is sourceless but satisfies the additional condition that the net flow through E_0 is 1, one needs to adapt the setting of the first sections in order to prove Theorem 5.2. For any $e \in \mathsf{E}_{1/2}^n$, let

$$\Theta_e^n := \{ \text{ unit flows } \theta \text{ from } e_- \text{ to } e_+ \text{ s.t. } \sum_{e' \in E_0} \theta(e') = \mathbb{1}_{e \in E_0}\}$$

From (13), one sees that for any unit flow θ from e_- to e_+, and any i,

$$\sum_{e' \in E_i} \theta(e') = \sum_{e' \in E_0} \theta(e') + \mathbb{1}_{e \in E_i} - \mathbb{1}_{e \in E_0} \ .$$

Thus, one sees that for any translation τ on the torus, and any edge e,

$$\theta \in \Theta_e^n \Leftrightarrow \theta \circ \tau \in \Theta_{\tau^{-1}(e)}^n \ . \tag{16}$$

Then, one may show as in the proof of Lemma 5.3 that $\theta \mapsto \mathcal{E}_r(\theta)$ has a unique minimizer on Θ_e^n, that we shall call $j_{r,n}^e$, and which is characterized by the same pseudo-Kirchhoff cycle law. The role of $j_{r,n}^e$ will be similar to that of i_r^e in the first sections, as hinted by the following lemma.

Lemma 5.4 *The functions* $r \mapsto i^{per}_{r,n}(e)$*, for any edge* e*, and* $r \mapsto \mathcal{R}_n(r)$ *admit partial derivatives of all order. In addition, for any edges* e*,* e'*,*

(i) $\forall e' \neq e, \quad \partial_{e'} i^{per}_{r,n}(e) = \frac{i^{per}_{r,n}(e')}{r(e')} j^{e'}_{r,n}(e) = \frac{i^{per}_{r,n}(e')}{r(e)} j^{e}_{r,n}(e')\ .$

(ii) $\forall e, \quad \partial_e i^{per}_{r,n}(e) = \frac{i^{per}_{r,n}(e)}{r(e)} (j^e_{r,n}(e) - 1)\ .$

(iii) $\forall e, \quad \partial_e \mathcal{R}_n(r) = (i^{per}_{r,n}(e))^2\ .$

Proof: The fact that $r \mapsto i^{per}_{r,n}(e)$ and $r \mapsto \mathcal{R}_n(r)$ admits partial derivatives of all order is analogous to the classical case, cf. the proof of Lemma 2.2.

Let us fix some edge $e' \in \mathsf{E}^n_{1/2}$. One may thus differentiate the node law and the pseudo-Kirchhoff cycle law of Lemma 5.3 with respect to $r(e')$. to obtain:

$$\forall x \in \mathsf{G}^{per}_n, \quad d^*[\partial_{e'} i^{per}_{r,n}](x) = 0\ ,$$

and

$$\forall h \in \overrightarrow{\Theta}, \quad \sum_e h(e) r(e) \left(\partial_{e'} i^{per}_{r,n}(e) + \frac{i^{per}_{r,n}(e')}{r(e')} \chi_{e'}(e) \right) = 0\ .$$

Thus, if we define:

$$\theta(e) = \partial_{e'} i^{per}_{r,n}(e) + \frac{i^{per}_{r,n}(e')}{r(e')} \chi_{e'}(e)\ ,$$

we see that:

$$\forall x \notin e', \quad d^*\theta(x) = 0\ ,$$

and,

$$\forall h \in \overrightarrow{\Theta}, \quad (h, \theta)_r = 0\ .$$

Furthermore,

$$d^*\theta(e'_-) = \frac{i^{per}_{r,n}(e')}{r(e')}\ ,$$

and:

$$\sum_{e \in E_0} \theta(e) = \frac{i^{per}_{r,n}(e')}{r(e')} \mathbb{1}_{e' \in E_0}\ .$$

It follows from the characterization of $j^{e'}_{r,n}$ that:

$$\theta = \frac{i^{per}_{r,n}(e')}{r(e')} j^{e'}_{r,n}\ .$$

This gives the proof of the first two equations. The proof of the last one is analogous to the classical case, cf. Lemma 2.2. □

When n is large, we would like to compare $j^e_{r,n}$ to a flow on the whole lattice $\mathbb{Z}^d$. To do this, we shall couple the network $(\mathbb{Z}^d, \mathbb{E}^d)$ with all the tori as follows. *This construction will be used throughout the section.*

We shall identify all the set of edges E_n^d, equipped with their resistances, as subsets of $\mathbb{E}^d$. First, fix e to be any edge such that e_- is the origin of $\mathbb{Z}^d$. Let $r \in [1, \Lambda]^{\mathbb{E}^d_{1/2}}$ be a fixed collection of resistances and define:

$$\mathsf{V}_n = \{-\lfloor n/2 \rfloor, \dots, \lfloor (n-1)/2 \rfloor\}^d$$

$$\mathsf{E}^n_{1/2} = \{(x,y) \in \mathsf{V}_n \times \mathbb{Z}^d \text{ s.t. } \exists i \in \{1, \dots, n\}\ y_i - x_i = 1 \text{ and } \forall j \neq i,\ x_j = y_j\} \ .$$

Now, we let G_n^{per} be the network with edge set $\mathsf{E}^n_{1/2}$, resistances induced by r and set of vertices all the endpoints of the edges of $\mathsf{E}^n_{1/2}$ with periodic condition, i.e vertices with identical coordinates modulo n are identified. Clearly, G_n^{per} is isomorphic to $\mathbb{T}^d_n$, and we shall thus use the notations E_0, Θ^n_e, Θ^n_0 and $j^e_{r,n}$ on G_n^{per} as well.

Lemma 5.5 *Let e be any fixed edge such that e_- is the origin of $\mathbb{Z}^d$. Let i^e_r be the minimal current on $\mathbb{Z}^d$ from e_- to e_+ , then $j^e_{r,n}$ converges to i^e_r in $\ell^2_-(\mathbb{E}^d, r)$.*

Proof: We let G_n^F be the network induced by the set of vertices V_n. Also, we define G_n^W be the network obtained from $\mathbb{Z}^d$ by identifying all the vertices outside $\{-\lfloor n/2 \rfloor + 1, \dots, \lfloor (n-1)/2 \rfloor\}^d$. See Figure 3.



G_n^{per} G_n^F G_n^W

Figure 3: The three graphs G_n^{per}, G_n^F and G_n^W when $n = 3$ and $d = 2$. Squares are identified as a single vertex, while empty circles are subject to periodic identification modulo n.

Now, let $\bigstar_n$ (resp. $\bigstar_n^{per}$) be the linear subspace of $\ell^2_-(\mathsf{E}^n_{1/2})$ spanned by the stars of G_n^W (resp. G_n^{per}) and $\diamondsuit_n$ (resp. $\diamondsuit_n^{per}$) the linear subspace spanned by the cycles of G_n^F (resp. G_n^{per}). Then, for any n,

$$\bigstar_n \subset \bigstar_n^{per} \qquad \text{and } \diamondsuit_n \subset \diamondsuit_n^{per} \cap \{\theta \in \ell^2_-(\mathsf{E}^n_{1/2}) \text{ s.t. } \sum_{e \in E_0} \theta(e) = 0\} \ .$$

If one defines:

$$\overrightarrow{\Theta}_n = \{\theta \in \diamondsuit_n^{per} \text{ s.t. } \sum_{e \in E_0} \theta(e) = 0\} \ ,$$

then $j^e_{r,n}$ is the orthogonal projection in $\ell^2_-(\mathsf{E}^n_{1/2}, r)$ of χ_e on $H_n = \overrightarrow{\Theta}_n^{\perp}$. Notice that it is a flow between e_- and e_+ on G_n^{per} but not necessarily on $\mathbb{Z}^d$. The inclusions above give:

$$\bigstar_n \subset H_n \subset \diamondsuit_n^{\perp} \ .$$

Let $\bigstar$ (resp. $\diamondsuit$) denote the closed linear subspace spanned by stars (resp. cycles) in $\ell^2_-(\mathbb{E}^d_{1/2}, r)$. Recall (cf. section 2.1), that $P_{\bigstar_n}\chi_e$ (resp. $P_{\diamondsuit_n^{\perp}}\chi_e$)

converges to $i_r^e = P_{\bigstar}\chi_e$, the minimal current on $\mathbb{Z}^d$ from e_- to e_+ (resp. $i_r^{F,e} = P_{\diamondsuit^\perp}\chi_e$, the free current on $\mathbb{Z}^d$ from e_- to e_+). But on $\mathbb{Z}^d$, currents are unique, and one other way to say it is to say that $\bigstar = \diamondsuit^\perp$. As a consequence, $(j_{r,n}^e)_{n\geq 1}$ converges also to i_r^e, the minimal current on $\mathbb{Z}^d$ from e_- to e_+. $\square$

In order to adapt the notion of graphs with homogeneous currents to this setting of sequences of finite graphs, define:

$$\alpha(d,L,\Lambda) := \sup_{n\geq 1} \sup_{r\in[1,\Lambda]^{\mathsf{E}_{1/2}}} \sup_{e\in\mathbb{T}_n^d} \sum_{d(e',e)\geq L} r(e')(j_{r,n}^e(e'))^2 \ .$$

Proposition 5.6 *The d-dimensional discrete tori have homogeneous currents in the sense that for any* $\Lambda \geq 1$,

$$\alpha(d,L,\Lambda) \xrightarrow[L\to+\infty]{} 0 \ .$$

Proof: We will adapt the proof of Proposition 3.9, using the convergence of $j_{r,n}^e$ given by Lemma 5.5. Let $\mathsf{V} = \mathbb{Z}^d$ and $\mathsf{E} = \mathbb{E}^d$.

Let $e_1, \ldots, e_d$ be the d edges going from 0 to some point of non-negative coordinates. Thanks to the translation property (16), one has, for any translation τ on the torus:

$$j_{r\circ\tau^{-1},n}^{\tau(e)} \circ \tau = j_{r,n}^e \ .$$

Thus,

$$\alpha(d,L,\Lambda) := \sup_{n\geq 1} \sup_{e\in\{e_1,\ldots,e_d\}} \sup_{r\in[1,\Lambda]^{\mathsf{E}_{1/2}}} \sum_{d(e',e)\geq L} r(e')(j_{r,n}^e(e'))^2 \ .$$

Let e be any fixed edge such that e_- is the origin, and define:

$$c := \limsup_{L\to\infty} \sup_{n\geq 1} \sup_{r\in[1,\Lambda]^{\mathsf{E}_{1/2}}} \sum_{d(e',e)\geq L} r(e')(j_{r,n}^e(e'))^2 \ .$$

It is thus enough to prove that $c = 0$. Notice that when n is fixed,

$$\limsup_{L\to\infty} \sup_{r\in[1,\Lambda]^{\mathsf{E}_{1/2}}} \sum_{d(e',e)\geq L} r(e')(j_{r,n}^e(e'))^2 = 0 \ .$$

Thus, one may find a sequence $(r_L, n_L)_{L\geq 1}$ in $[1,\Lambda]^{\mathsf{E}_{1/2}} \times \mathbb{N}$ such that:

$$c = \lim_{L\to\infty} \sum_{d(e',e)\geq L} r_L(e')(j_{r_L,n_L}^e(e'))^2 \ ,$$

and $n_L \xrightarrow[L\to\infty]{} +\infty$. The sequence j_{r_L,n_L}^e is bounded in $\ell^2_-(\mathsf{E}_{1/2})$. Thus, by compactness of $[1,\Lambda]^{\mathsf{E}_{1/2}}$ and weak compactness of balls in $\ell^2_-(\mathsf{E}_{1/2})$, one may extract a sequence from $(r_L, n_L)_{L\geq 1}$, that we shall still denote by $(r_L, n_L)_{L\geq 1}$ to lighten the notations, such that $(j_{r_L,n_L}^e)_L$ converges weakly to some $\theta \in \ell^2_-(\mathsf{E}_{1/2})$, and $(r_L)_L$ converges to some $r \in [1,\Lambda]^{\mathsf{E}_{1/2}}$. Notice that θ is a flow on the whole lattice $\mathbb{Z}^d$ since for any n, $(j^e_{r_L,n_L})_L$ is a unit flow on G_n^{per}. Using the minimality of i^e_r,

$$\begin{aligned}
\mathcal{E}_r(i^e_r) &\leq \mathcal{E}_r(\theta)\ , \\
&= \sum_{e'\in \mathsf{E}_{1/2}} r(e')\theta^2(e')\ , \\
&\leq \limsup_{L\to\infty} \left(\sum_{d(e',e)\leq L} r_L(e')\left(j^e_{r_L,n_L}(e')\right)^2 \right)\ , \\
&= \limsup_{L\to\infty} \left(\mathcal{E}_{r_L}(j^e_{r_L,n_L}) - \sum_{d(e',e)>L} r_L(e')\left(j^e_{r_L,n_L}(e')\right)^2 \right)\ , \\
&\leq \limsup_{L\to\infty} \mathcal{E}_{r_L}(j^e_{r_L,n_L}) - c\ , \\
&\leq \limsup_{L\to\infty} \mathcal{E}_{r_L}(j^e_{r,n_L}) - c\ , \qquad (17)
\end{aligned}$$

where in the last inequality we used the minimality property of $j^e_{r_L,n_L}$. Now, using Minkowski's inequality,

$$\begin{aligned}
\mathcal{E}_{r_L}(j^e_{r,n_L}) &= \sum_{e'\in \mathsf{E}_{1/2}} r_L(e')(j^e_{r,n_L}(e'))^2\ , \\
&\leq \left(\sqrt{\sum_{e'\in \mathsf{E}_{1/2}} r_L(e')(i^e_r(e'))^2} + \sqrt{\Lambda \sum_{e'\in \mathsf{E}_{1/2}} (j^e_{r,n_L}(e') - i^e_r(e'))^2} \right)^2 .
\end{aligned}$$

Since $(r_L)_{L\geq 1}$ converges simply to r and is bounded by Λ, the dominated convergence theorem gives:

$$\sum_{e'\in \mathsf{E}_{1/2}} r_L(e')(i^e_r(e'))^2 \xrightarrow[L\to\infty]{} \mathcal{E}_r(i^e_r)\ ,$$

Lemma 5.5 says that $(j^e_{r,n_L})_{L\geq 1}$ converges to i^e_r in $\ell^2_-(\mathsf{E}_{1/2})$. Thus:

$$\limsup_{L\to\infty} \mathcal{E}_{r_L}(j^e_{r,n_L}) \leq \mathcal{E}_r(i^e_r)\ .$$

Plugging this into (17) shows that $c = 0$. ☐

Now, one may complete the proof of Theorem 5.2.

Proof: **of Theorem 5.2.** With Lemma 5.4 at hand, it is easy to reproduce the proofs of Lemma 2.3, Corollary 3.12 and Theorem 3.11 with $\alpha(\mathsf{G}, L, \Lambda)$ replaced by $\alpha(d, L, \Lambda)$. One obtains notably the existence of constants $C(\Lambda)$ and $C(\Lambda, d)$ such that for any n, and any $L \geq 1$,

$$\sum_{diam(S)\geq L} \|(\mathcal{R}_n)_S\|_2^2 \leq C(\Lambda)\alpha(G, L, \Lambda) \sum_{S\neq\emptyset} \|(\mathcal{R}_n)_S\|_2^2\ ,$$

and

$$\sum_e \|(\mathcal{R}_n)_{\{e\}}\|_2^2 \leq \mathsf{Var}(\mathcal{R}_n) = \sum_{S\neq\emptyset} \|(\mathcal{R}_n)_S\|_2^2 \leq C(\Lambda, d) \sum_e \|(\mathcal{R}_n)_{\{e\}}\|_2^2\ .$$

Now, thanks to the translation invariance of the model given by (14) and the fact that the edge-resistances are i.i.d,

$$\beta_n := \sup_n \sup_{e \in \mathsf{E}^n_{1/2}} \frac{\|(\mathcal{R}_n)_{\{e\}}\|_2^2}{\mathsf{Var}(\mathcal{R}_n)} = \Theta(1/n^d) \ ,$$

and,

$$\sup_{e \in \mathsf{E}^n_{1/2}} \mathbb{E}[(i^{per}_{r,n}(e))^2] = \Theta(\frac{1}{n^d}\mathbb{E}(\mathcal{R}_n)) = \Theta(\frac{1}{n^{2d-2}}) \ .$$

This already shows that:

$$\mathsf{Var}(\mathcal{R}_n) = \Theta(\sup_{e \in \mathsf{E}^n_{1/2}} \mathbb{E}[(i^{per}_{r,n}(e))^2]^2) = \Theta(\frac{1}{n^{3d-4}}) \ .$$

Now, Proposition 5.6 allows to repeat the proof of Theorem 4.1. Let F_n be the distribution function of $\frac{\mathcal{R}_n - \mathbb{E}(\mathcal{R}_n)}{\sqrt{\mathsf{Var}(\mathcal{R}_n)}}$. One obtains the existence of a function f having limit 0 at 0^+ and such that for any n,

$$\|F_n - \Phi\|_\infty \leq f(\beta_n) \ .$$

This concludes the proof of the central limit theorem. □

6 Perspectives

We end this article with some questions left open.

First, it is not clear whether the notion of homogeneous currents is really useful to get a central limit theorem. For instance, in the counterexample of Figure 1, one sees that the currents i^e_r still spread most of their mass at very localized places, namely near e and near the edge e'_k which is in the same connected component as e. Thus, one should be able to adapt the proof of the central limit theorem in this special case. One may conjecture that only bounded degree and small influences are enough to get a central limit theorem. On the other hand, if the homogeneous currents hypothesis is proved really necessary, it would be important to understand which graphs satisfy it, and whether it is stable under perturbations like quasi-isometries.

Second, the most obvious question left open is the one raised in the introduction, that is to determine the order of the variance and to show a central limit theorem for the resistance on the cube of side length n in $\mathbb{Z}^d$, and not only on the torus. More generally, consider a domain Ω of $\mathbb{R}^d$ with two disjoint subsets of its boundary, A and Z. Let G_n be the graph induced by $\Omega \cap \frac{1}{n}\mathbb{Z}^d$ and let $\mathcal{R}_n$ be the effective resistance between A and Z on G_n. Then, we conjecture that a Gaussian central limit theorem holds for $\mathcal{R}_n$.

7 Acknowledgements

I would like to thank Itai Benjamini and Michel Benaim with whom I began this research. Thanks also to Pierre Mathieu, Alano Ancona, Daniel Boivin and

Thierry Delmotte for very useful discussions. Last but not least, the main part of Proposition 3.9 is due to Mickaël de la Salle, I owe a lot to him. This work was partially supported by the ANR project GeMeCoD ANR 2011 BS01 007 01.

A Proofs of section 3.1

A.1 Proof of Proposition 3.1

Let us first suppose that $\mathsf{E}_{1/2}$ is finite. Then, Proposition 3.1 is well known, cf. [6], but we shall quickly recall the proof for the sake of completeness.

For a subset $S \subset \mathsf{E}_{1/2}$, let L_S be the operator on $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$ defined by:

$$L_S f(r) = \int f(r) \prod_{e \in S} d\mu_e(r(e)) \ .$$

Let 1 denote the identity operator. Notice that $L_{\{e\}}$ and $L_{\{e'\}}$ commute for any e and e'. Since $\Delta_{\{e'\}} = 1 - L_{\{e\}}$,

$$1 = \prod_{e \in E} (\Delta_{\{e\}} + L_{\{e\}}) = \sum_{S \subset E} L_{S^c} \prod_{e \in S} \Delta_{\{e\}} \ .$$

Since

$$f_S = \sum_{S \subset E} L_{S^c} \prod_{e \in S} \Delta_{\{e\}} f \ ,$$

this shows that $f_\emptyset = \mathsf{E}_{1/2}E(f)$ and $f = \sum_{S \subset \mathsf{E}_{1/2}} f_S$. Now, remark that for any e,

$$L_e \Delta_e = 0 \ .$$

Thus, for any S and any $e \in S$,

$$L_e f_S = 0 \ .$$

This implies that $\Delta_e f = \sum_{S \ni e} f_S$. Now, if $S \neq S'$, suppose for instance that there is some $e \in S \setminus S'$:

$$\mathsf{E}_{1/2}E[f_S g_{S'}] = \text{Ł}_{\mathsf{E}_{1/2}}(f_S g_{S'}) = \text{Ł}_{\mathsf{E}_{1/2}} L_{\{e\}}(f_S g_{S'}) = \text{Ł}_{\mathsf{E}_{1/2}}(g_{S'} L_{\{e\}}(f_S)) = 0 \ .$$

Now, let us suppose that $\mathsf{E}_{1/2}$ is countable, and fix some exhaustion E_n of the edges: E_n is finite for any n, $E_n \subset E_{n+1}$ and $\mathsf{E}_{1/2} = \cup_n E_n$. Denote by f_n the conditional expectation of f with respect to r_{E_n}. We have:

$$f_n = L_{E_n^c} f$$

and thus,

$$(f_n)_S = f_S \mathbb{1}_{S \subset E_n} \ , \tag{18}$$

which implies:

$$f_n = \sum_{S \subset E_n} f_S \text{ and } \|f_n\|_2^2 = \sum_{S \subset E_n} \|f_S\|_2^2 \ .$$

Since $(f_n)_{n \in \mathbb{N}}$ converges to f in L^2, we know that $\|f_n\|_2^2$ converges to $\|f\|_2^2$. Thus, $\sum_S f_S$ converges normally to f in L^2 and for any S. All the other properties can then be derived from (18).

A.2 Proof of Proposition 3.3

Let r and r' be two independent copies of law $\mathbb{P}$. Let S_ε be the (possibliy infinite) random subset of E drawn at random as follows: $(\mathbb{1}_{e\in S_\varepsilon})_{e\in\mathsf{E}_{1/2}}$ are i.i.d with distribution Bernoulli of parameter $\varepsilon \in [0,1]$, independent of (r,r'). Now, we define the following linear operators from $L^2(\mathbb{R}^{\mathsf{E}_{1/2}})$ to $L^2(\mathbb{R}^{\mathsf{E}_{1/2}} \times \mathbb{R}^{\mathsf{E}_{1/2}})$:

$$L^{r'}_{\{e\}} f(r) = f(r^{e\leftarrow r'}) \ .$$

Then, the noisy version of $f(r)$ may be written as:

$$f(r^\varepsilon) = \prod_{e\in S_\varepsilon} L^{r'}_{\{e\}} f(r) \ .$$

Notice that:

$$\begin{cases} \forall e \neq e', & L^{r'}_{\{e\}} L_{\{e'\}} & = & L_{\{e'\}} L^{r'}_{\{e\}} \\ \forall e, & L^{r'}_{\{e\}} L_{\{e\}} & = & L_{\{e\}} \ , \\ \forall e, & L_{\{e\}} L^{r'}_{\{e\}} & = & L^{r'}_{\{e\}} \ . \end{cases}$$

Thus,

$$L^{r'}_{\{e\}} \Delta_e = L^{r'}_{\{e\}} - L_{\{e\}} \ .$$

Whence:

$$\begin{aligned} f(r^\varepsilon) &= \sum_{S\subset\mathsf{E}_{1/2}} \prod_{e\in S_\varepsilon} L^{r'}_{\{e\}} f_S(r) \ , \\ &= \sum_{S\subset\mathsf{E}_{1/2}} \prod_{e\in S_\varepsilon} L^{r'}_{\{e\}} \prod_{e\in S^c} L_{\{e\}} \prod_{e\in S} \Delta_e f(r) \ , \\ &= \sum_{S\subset\mathsf{E}_{1/2}} \prod_{e\in S^c} L_{\{e\}} \prod_{e\in S\cap S_\varepsilon} (L^{r'}_{\{e\}} - L_{\{e\}}) \prod_{e\in S\setminus S_\varepsilon} \Delta_e f(r) \ . \end{aligned}$$

Now, denote by $L'_{\{e\}}$ the operator on $L^2(\mathbb{R}^{\mathsf{E}_{1/2}} \times \mathbb{R}^{\mathsf{E}_{1/2}})$ which integrates $r'(e)$. Notice that:

$$L'_{\{e\}}(L^{r'}_{\{e\}} - L_{\{e\}}) = 0 \text{ and } L'_{\{e\}}\Delta_e = \Delta_e \ .$$

Then,

$$\begin{aligned} \mathbb{E}[f(r^\varepsilon)|r, S_\varepsilon] &= \prod_{e\in\mathsf{E}_{1/2}} L'_{\{e\}}((r,r') \mapsto f(r^\varepsilon)) \ , \\ &= \sum_{S\subset\mathsf{E}_{1/2}} \prod_{e\in S^c} L_{\{e\}} \prod_{e\in S\setminus S_\varepsilon} \Delta_e f(r) \mathbb{1}_{S\cap S_\varepsilon=\emptyset} \ , \\ &= \sum_{S\subset\mathsf{E}_{1/2}} \prod_{e\in S^c} L_{\{e\}} \prod_{e\in S} \Delta_e f(r) \mathbb{1}_{S\cap S_\varepsilon=\emptyset} \ . \end{aligned}$$

Thus,

$$\mathbb{E}[f(r^\varepsilon)|r] = \sum_{S\subset\mathsf{E}_{1/2}} f_S(r)\mathbb{P}(S\cap S_\varepsilon = \emptyset) = \sum_{S\subset\mathsf{E}_{1/2}} (1-\varepsilon)^{|S|} f_S(r) \ .$$

References

[1] M. Benaïm and R. Rossignol. Exponential concentration for first passage percolation through modified Poincaré inequalities. *Ann. Inst. Henri Poincaré Probab. Stat.*, 44(3):544–573, 2008.

[2] I. Benjamini, G. Kalai, and O. Schramm. Noise sensitivity of Boolean functions and applications to percolation. *Publ. Math. I.H.E.S.*, 90:5–43, 1999.

[3] D. Boivin. Tail estimates for homogenization theorems in random media. *ESAIM Probab. Stat.*, 13:51–69, 2009.

[4] D. Boivin and J. Depauw. Spectral homogenization of reversible random walks on $\mathbb{Z}^d$ in a random environment. *Stochastic Process. Appl.*, 104(1):29–56, 2003.

[5] B. Bollobás. *Modern graph theory.* Springer, 1998.

[6] J. Bourgain. *Walsh subspaces of L^p-product spaces*, pages IV.1–IV.9. Séminaire d'analyse fonctionnelle de l'école Polytechnique. Palaiseau, 1979.

[7] S. Chatterjee. Chaos, concentration, and multiple valleys. *ArXiv e-prints*, October 2008.

[8] L. H. Y. Chen and Q-M. Shao. Normal approximation under local dependence. *Ann. Probab.*, 32(3A):1985–2028, 2004.

[9] T. Delmotte. Inégalité de Harnack elliptique sur les graphes. *Colloq. Math.*, 72(1):19–37, 1997.

[10] T. Delmotte and J.-D. Deuschel. On estimating the derivatives of symmetric diffusions in stationary random environment, with applications to $\nabla\phi$ interface model. *Probab. Theory Related Fields*, 133(3):358–390, 2005.

[11] B. Efron and C. Stein. The jackknife estimate of variance. *Ann. Statist.*, 9(3):586–596, 1981.

[12] A. Gloria and F. Otto. An optimal variance estimate in stochastic homogenization of discrete elliptic equations. *Ann. Probab.*, 39(3):779–856, 2011.

[13] H. Hatami. A structure theorem for Boolean functions with small total influences. *Annals of math.*, 176(1):509–533, 2012.

[14] V. V. Jikov, S. M. Kozlov, and O. A. Oleinik. *Homogenization of differential operators and integral functionals.* Springer-Verlag, 1994.

[15] G. Kirchhoff. On the solution of the equations obtained from the investigation of the linear distribution of galvanic currents. *IRE Transactions on circuit theory*, 5(1), 1958.

[16] S. M. Kozlov. Average difference schemes. *Mat. Sb. (N.S.)*, 129(171)(3):338–357, 447, 1986.

[17] R. Künnemann. The diffusion limit for reversible jump processes on $\mathbb{Z}^d$ with ergodic random bond conductivities. *Comm. Math. Phys.*, 90(1):27–68, 1983.

[18] R. Lyons and Y. Peres. *Probability on trees and networks.* Cambridge University Press, 2012. In preparation. Current version available at `http://mypage.iu.edu/~rdlyons/`.

[19] J. Moser. On Harnack's theorem for elliptic differential equations. *Comm. Pure Appl. Math.*, 14:577–591, 1961.

[20] E. Mossel. Gaussian bounds for noise correlation of functions. *Geom. Funct. Anal.*, 19(6):1713–1756, 2010.

[21] A. Naddaf and T. Spencer. Estimates on the variance of some homogenization problems. Unpublished., 1998.

[22] G. Papanicolaou and S. Varadhan. Boundary value problems with rapidly oscillating random coefficients. In *Random fields, Vol. I, II (Esztergom, 1979)*, volume 27 of *Colloq. Math. Soc. János Bolyai*, pages 835–873. North-Holland, Amsterdam, 1981.

[23] J. Wehr. A lower bound on the variance of conductance in random resistor networks. *J. Statist. Phys.*, 86(5-6):1359–1365, 1997.